

Anomaly freedom in Seiberg-Witten noncommutative gauge theories

Friedemann Brandt ^{a 1}, C.P. Martín ^{b 2} and F. Ruiz Ruiz ^{c 3}

^a *Max-Planck-Institute for Mathematics in the Sciences,
Inselstraße 22-26, D-04103 Leipzig, Germany*

^{b, c} *Departamento de Física Teórica I, Facultad de Ciencias Físicas
Universidad Complutense de Madrid, 28040 Madrid, Spain*

We show that noncommutative gauge theories with arbitrary compact gauge group defined by means of the Seiberg-Witten map have the same one-loop anomalies as their commutative counterparts. This is done in two steps. By explicitly calculating the $\epsilon^{\mu_1\mu_2\mu_3\mu_4}$ part of the renormalized effective action, we first find the would-be one-loop anomaly of the theory to all orders in the noncommutativity parameter $\theta^{\mu\nu}$. And secondly we isolate in the would-be anomaly radiative corrections which are not BRS trivial. This gives as the only true anomaly occurring in the theory the standard Bardeen anomaly of commutative space-time, which is set to zero by the usual anomaly cancellation condition.

Keywords: anomaly freedom, Seiberg-Witten map, noncommutative gauge theories.

¹E-mail: fbrandt@mis.mpg.de

²E-mail: carmelo@elbereth.fis.ucm.es

³E-mail: t63@aeneas.fis.ucm.es

1 Introduction

It is a well known fact that not all relevant gauge groups in particle physics are consistent with the Moyal product of noncommutative field theory. An example of this is provided by the Moyal product $A_\mu(x) \star A_\nu(x)$ of two $SU(N)$ Lie algebra valued gauge fields $A_\mu(x)$ and $A_\nu(x)$. It is clear that such product does not lie in the $SU(N)$ Lie algebra but in a representation of its enveloping algebra, so $A_\mu(x)$ can not be regarded as a truly noncommutative $SU(N)$ gauge field. This makes it difficult to formulate, even classically, noncommutative extensions of some physically relevant gauge theories like *e.g.* the standard model. A way to circumvent this problem [1, 2] is to build noncommutative gauge and matter fields from ordinary ones by means of the Seiberg-Witten map [3]. Using this approach, classical noncommutative gauge theories have been constructed for arbitrary compact groups [1, 2, 4, 5] and noncommutative gauge theories with $SU(5)$ and $SO(10)$ gauge groups have been constructed in ref. [13]. Furthermore, a noncommutative standard model has been formulated in ref. [6] and some of its phenomenological consequences have been explored in a number of papers [7, 8, 9, 10, 11, 12]. Many of these noncommutative gauge theories, among them the noncommutative standard model, involve chiral fermions, so the corresponding classical gauge symmetry may be broken by quantum corrections. In other words, an anomaly may occur and the resulting quantum theory may then become inconsistent. To study the consistency of quantum noncommutative gauge theories defined by means of the Seiberg-Witten map, it is therefore necessary to study whether new types of anomalies occur *-i.e.* anomalies which do not appear in ordinary commutative spacetime and hence that may require additional anomaly cancellation conditions.

In refs. [14, 15, 16] it has been shown that for Yang-Mills type gauge theories with arbitrary semisimple gauge groups the only nontrivial solution to the anomaly consistency condition is the usual Bardeen anomaly, regardless of whether or not the theory is Lorentz invariant or renormalizable by power counting. This result readily applies to gauge noncommutative field theories constructed by means of the Witten-Seiberg map, since, as far as these matters are concerned, the presence of a noncommutative matrix parameter $\theta^{\mu\nu}$ with mass dimension -2 only precludes Lorentz invariance and power-counting renormalizability. Thus, for noncommutative gauge theories with semisimple gauge groups, there are no $\theta^{\mu\nu}$ -dependent anomalies and any $\theta^{\mu\nu}$ -dependent breaking of the BRS identity, being cohomologically trivial, can be set to zero by adding appropriate counterterms to the effective action. Note that the addition of these $\theta^{\mu\nu}$ -dependent counterterms to the effective action makes sense within the framework of

effective field theory, but this agrees with the observation that noncommutative field theories defined by means of the Seiberg-Witten map should be considered as effective field theories [13, 17]. All the above implies that no anomalous $\theta^{\mu\nu}$ -dependent terms should occur in the Green functions of noncommutative theories with semisimple gauge groups, a fact that has been proved to hold true at order one in $\theta^{\mu\nu}$ for the three-point function of the gauge field and a simple gauge group by explicit computation of the appropriate Feynman diagrams [18].

The situation is very different if the gauge group is not semisimple. In this case, the consistency condition for gauge anomalies has other nontrivial solutions besides Bardeen's anomaly. In particular, in four dimensions and if the gauge group is $G \times U(1)_Y$, with G a semisimple gauge group, the additional nontrivial solutions are of the form

$$\int d^4x \, c \, \text{I}_{\text{inv}}[f_{\mu\nu}, G_{\mu\nu}] . \quad (1.1)$$

Here matter fields have been integrated out, c is the $U(1)_Y$ ghost field and $\text{I}_{\text{inv}}[f_{\mu\nu}, G_{\mu\nu}]$ is a gauge invariant function of the $U(1)_Y$ field strength $f_{\mu\nu}$, the G field strength $G_{\mu\nu}$ and their covariant derivatives. Note that there are infinitely many candidate anomalies of this type since neither power counting nor Lorentz invariance are available to reduce the number of invariants $\text{I}_{\text{inv}}[f_{\mu\nu}, G_{\mu\nu}]$. Furthermore, when the gauge group contains more than one abelian factor, there are additional candidate anomalies of yet another type [14, 15, 16]. The purpose of this paper is to investigate whether anomalies of these types occur in noncommutative gauge theories with nonsemisimple gauge groups defined through the Seiberg-Witten map. This is not a trivial question and has far reaching implications. Indeed, did solutions of type (1.1) occur in perturbation theory, the corresponding quantum gauge theory would be anomalous, the anomaly being $\theta^{\mu\nu}$ -dependent. To remove the resulting anomaly and render the quantum theory consistent, one would then have to impose constraints on the fermions hypercharges. A conspicuous instance of a model with such a gauge group for which this point should be cleared is the noncommutative standard model [6].

In this paper we will prove that, for a noncommutative field theory with arbitrary compact gauge group defined by means of the Seiberg-Witten map, the only anomaly that occurs at one loop (hence, to all orders in perturbation theory, if one assumes the existence of a nonrenormalization theorem for the anomaly) is the usual Bardeen anomaly. The paper is organized as follows. In Section 2 we fix the notation, define the chiral BRS transformations and use the Seiberg-Witten map to classically define the noncommutative model. Section 3 uses dimensional regularization to explicitly compute the $\epsilon^{\mu_1\mu_2\mu_3\mu_4}$ part of the renormalized effective action. This yields a complicated power series in the noncommutativity parameter

$\theta^{\mu\nu}$, of which the term of order zero is the usual Bardeen anomaly of commutative field theory. In Section 4 we show that all terms in this series of order one or higher in $\theta^{\mu\nu}$ are cohomologically trivial with respect to the chiral BRS operator and find the counterterm that removes them from the renormalized effective action. Section 5 contains our conclusions. We postpone to two appendices some very technical points of our arguments. Let us emphasize that in this paper we will only discuss gauge anomalies –see refs. [19, 20] for related work on the rigid axial anomaly.

2 The model, notation and conventions

Let us consider a compact nonsemisimple gauge group $G = G_1 \times \cdots \times G_N$, with G_i a simple compact group if $i = 1, \dots, s$ and an abelian group if $i = s+1, \dots, N$. We may assume without loss of generality that the abelian factors come with irreducible representations, which of course are one-dimensional. Let us denote by $\psi_{i_1 \dots i_s}$ a Dirac field on ordinary Minkowski spacetime carrying an arbitrary unitary irreducible representation of the Lie algebra of G . Since the abelian factors come with one-dimensional representations, the indices in the Dirac field $\psi_{i_1 \dots i_s}$ correspond to the simple factors. From now on we will collectively denote the “simple” indices $(i_1 \dots i_s)$ by the multi-index I . The corresponding vector potential v_μ on Minkowski spacetime in the representation of the Lie algebra carried by ψ_I will have the following decomposition in terms of the gauge fields a_μ^k and a_μ^l associated to the factors of the group G

$$v_\mu = \sum_{k=1}^s g_k (a_\mu^k)^a (T^k)^a + \sum_{l=s+1}^N g_l a_\mu^l T^l .$$

Here g_k and g_l are the coupling constants and $\{(T^k)^a, T^l\}$, with $a = 1, \dots, \dim G_k$ for every $k = 1, \dots, s$ and $l = s+1, \dots, N$, stand for the generators of the G Lie algebra in the unitary irreducible representation under consideration. As usual, a sum over a is understood. The matrix elements IJ of these generators are always of the form

$$(T^k)_{IJ}^a = \delta_{i_1 j_1} \cdots (T^k)_{i_k j_k}^a \cdots \delta_{i_s j_s} \\ T_{IJ}^l = \delta_{i_1 j_1} \cdots \delta_{i_s j_s} Y^l ,$$

where $(T^k)_{i_k j_k}^a$ are the matrix elements of the generator $(T^k)^a$ of the Lie algebra of the factor G_k in some given irreducible representation. Given any two generators $(T^k)_{IJ}^a$ and $(T^{k'})_{IJ}^{a'}$

as above we define the trace operation $\mathbf{Tr}$ as

$$\begin{aligned}\mathbf{Tr} (T^k)^a (T^{k'})^{a'} &= (T^k)_{IJ}^a (T^{k'})_{JI}^{a'} \\ &= \delta_{i_1 j_1} \cdots (T^k)_{i_k j_k}^a \cdots \delta_{i_s j_s} \delta_{j_1 i_1} \cdots (T^{k'})_{j_{k'} i_{k'}}^{a'} \cdots \delta_{j_s i_s} .\end{aligned}$$

The ghost field λ associated to v_μ , also in the representation furnished by ψ_I , is

$$\lambda = \sum_{k=1}^s g_k (\lambda^k)^a (T^k)^a + \sum_{l=s+1}^N g_l \lambda^l T^l ,$$

with $(\lambda^k)^a$ and λ^l being the ghost fields for the factors in G . Now we consider the theory that arises from chirally coupling, say left-handedly, the fermion field ψ_I to the gauge field v_μ . The fermionic part of the corresponding classical action reads

$$S^{\text{fermion}} = \int d^4x \bar{\psi}_I i \hat{D}(v)_{IJ} \psi_J , \quad (2.1)$$

with $\bar{\psi}_I = \psi_I^\dagger \gamma^0$ and

$$\hat{D}(v)_{IJ} \psi_J = \delta_{IJ} \not{\partial} \psi_J + \not{v}_{IJ} P_- \psi_J .$$

Here P_- is the left-handed chiral projector, given by

$$P_\pm = \frac{1}{2} (1 \pm \gamma_5) \quad \gamma_5 = -i \gamma^0 \gamma^1 \gamma^2 \gamma^3 ,$$

the gamma matrices γ^μ being defined by $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu}$ and the convention for the Minkowski metric $\eta_{\mu\nu}$ being $\eta_{\mu\nu} = \text{diag}(+, -, -, -)$. This action is invariant under the chiral BRS transformations

$$s v_\mu = \partial_\mu \lambda + [v_\mu, \lambda] \quad s \psi = -\lambda P_- \psi \quad s \bar{\psi} = \bar{\psi} \lambda P_+ \quad s \lambda = -\lambda \lambda . \quad (2.2)$$

As usual, the BRS operator s commutes with ∂_μ , satisfies the anti-Leibniz rule and is nilpotent, *i.e.* $s^2 = 0$.

To construct the noncommutative extension of the ordinary gauge theory defined by the classical action S^{fermion} , we use the formalism developed in refs. [1, 2, 4, 5]. To this end, we first define the noncommutative gauge field V_μ , the noncommutative spinor field Ψ_I and the noncommutative ghost field Λ in terms of their ordinary counterparts v_μ , ψ_I and λ by using the Seiberg-Witten map [3]. This is done as follows. The fields $V_\mu = V_\mu[v; \theta]$, $\Psi_I = \Psi_I[\psi, v; \theta]$ and $\Lambda = \Lambda[\lambda, v; \theta]$ are formal power series in $\theta^{\mu\nu}$, with coefficients depending on the ordinary fields and their derivatives, that take values in the representation of the enveloping algebra

of the Lie algebra of the group G furnished by the ordinary Dirac field ψ_I and solve the Seiberg-Witten equations

$$s_\star V_\mu = sV_\mu \quad s_\star \Psi = s\Psi \quad s_\star \Lambda = s\Lambda \quad (2.3)$$

subject to the boundary conditions

$$V_\mu[v; \theta=0] = v_\mu \quad \Psi_I[v, \psi; \theta=0] = \psi_I \quad \Lambda[\lambda, v; \theta=0] = \lambda . \quad (2.4)$$

In eq. (2.3) s is the ordinary BRS operator of eq. (2.2), while $s_\star$ denotes the noncommutative BRS chiral operator, whose action on the noncommutative fields is given by

$$s_\star V_\mu = \partial_\mu \Lambda + [V_\mu, \Lambda]_\star \quad s_\star \Psi = -\Lambda \star P_- \Psi \quad s_\star \Lambda = -\Lambda \star \Lambda . \quad (2.5)$$

The commutator $[f, g]_\star$ stands for

$$[f, g]_\star = f \star g - g \star f ,$$

with $f \star g$ the Moyal product of functions on Minkowski spacetime, defined for arbitrary f and g by

$$(f \star g)(x) = \int \frac{d^4 p}{(2\pi)^4} \int \frac{d^4 q}{(2\pi)^4} e^{-i(p+q)x} e^{-\frac{i}{2} \theta^{\alpha\beta} p_\alpha q_\beta} \tilde{f}(p) \tilde{g}(q) ,$$

$\tilde{f}(p)$ and $\tilde{g}(q)$ being the Fourier transforms of f and g . For the noncommutative field Ψ_I we further demand it to be linear in ψ_I , so that

$$\Psi_{\alpha I} = \left(\delta_{IJ} \delta_{\alpha\beta} + M[v, \partial, \gamma, \gamma_5; \theta]_{\alpha\beta IJ} \right) \psi_{\beta J} , \quad (2.6)$$

where α and β are Dirac indices. Note that, in accordance with the boundary condition for $\Psi_I[v, \psi; \theta]$, the differential operator $M[v, \partial, \gamma, \gamma_5; \theta]_{\alpha\beta IJ}$ vanishes at $\theta = 0$. Taking Ψ_i linear in ψ_I , as in eq. (2.6), is always possible [21] and is the natural choice within the framework of noncommutative geometry [17]. Once the noncommutative fields have been defined, one considers the following noncommutative classical action

$$S_{\text{nc}}^{\text{fermion}} = \int d^4 x \bar{\Psi}_I \star i \hat{D}(V)_{IJ} \Psi_J , \quad (2.7)$$

where $\bar{\Psi}_I = \bar{\Psi}_I^\dagger \gamma^0$ and

$$\hat{D}(V)_{IJ} \Psi_J = \delta_{IJ} \not{\partial} \Psi_J + V_{IJ} \star P_- \Psi_J .$$

We stress that the noncommutative fields are functions of ordinary fields as given by the Seiberg-Witten map and hence the noncommutative action is also a functional of these. Furthermore, the noncommutative action $S_{\text{nc}}^{\text{fermion}}$ is invariant under the ordinary chiral BRS transformations in eq. (2.2) since, by definition of the Seiberg-Witten map,

$$s S_{\text{nc}}^{\text{fermion}} = s_{\star} S_{\text{nc}}^{\text{fermion}}$$

and, by construction,

$$s_{\star} S_{\text{nc}}^{\text{fermion}} = 0 .$$

The effective action $\Gamma[v; \theta]$ of the noncommutative theory is formally defined by

$$\begin{aligned} \Gamma[v; \theta] &= -i \ln Z[v; \theta] \\ Z[v; \theta] &= \mathcal{N} \int [d\bar{\psi}] [d\psi] \exp \left(i S_{\text{nc}}^{\text{fermion}} \right) , \end{aligned} \tag{2.8}$$

with $\mathcal{N}$ a normalization constant chosen so that $Z[v=0; \theta] = 1$, *i.e.*

$$\mathcal{N}^{-1} = \int [d\bar{\psi}] [d\psi] \exp \left(\int d^4x \bar{\psi} i \not{\partial} \psi \right) ,$$

and $[d\bar{\psi}] [d\psi]$ the measure for ordinary fermion fields. Also formally, the invariance of $S_{\text{nc}}^{\text{fermion}}$ under s leads to the invariance of $\Gamma[v; \theta]$ under ordinary gauge transformations of v_{μ} . The problem is that all this is formal since defining the effective action requires renormalization. The question that should really be addressed is whether it is possible to define a renormalized effective action $\Gamma^{\text{ren}}[v; \theta]$ invariant under s . Were this the case, the theory would be anomaly free. In this paper we provide an answer in the negative and show that the anomaly has the same form as for the ordinary, *i.e.* commutative theory.

3 Form of the noncommutative anomaly

In this section we use dimensional regularization to define a renormalized effective action $\Gamma^{\text{ren}}[v; \theta]$ and find a closed expression for the anomaly $s\Gamma^{\text{ren}}[v; \theta]$ in terms of the noncommutative fields V_{μ} and Λ . To dimensionally regularize the theory, we consider the action

$$S_{\text{nc}}^{\text{reg}} = \int d^{2\omega}x \bar{\Psi}_I \star i \left(\delta_{IJ} \not{\partial} \Psi_J + \bar{\gamma}^{\mu} V_{\mu IJ} \star P_{-} \Psi_J \right) , \tag{3.1}$$

first introduced in the context of noncommutative gauge theories in ref. [18] for $U(N)$ theories and theories with simple groups. Here we use dimensional regularization *à la* Breitenlohner

and Maison [22]. We will use the notation in that reference, in which 4-dimensional objects are denoted with bars ($\bar{g}^\mu_\mu = 4$) and evanescent or $(2\omega-4)$ -dimensional quantities are denoted with hats ($\hat{g}^\mu_\mu = 2\omega-4$). The dimensionally regularized partition function $Z^{\text{reg}}[v; \theta]$ is defined as the sum of the dimensionally regularized Feynman diagrams generated by the path integral

$$Z^{\text{reg}}[v; \theta] = \mathcal{N} \int [d\bar{\psi}] [d\psi] \exp \left(i S_{\text{nc}}^{\text{reg}} \right) . \quad (3.2)$$

In the regularized partition function we perform the change of variables $\psi_{\beta J}, \bar{\psi}_{\beta J} \rightarrow \Psi_{\alpha I}, \bar{\Psi}_{\alpha I}$, with

$$\begin{aligned} \Psi_{\alpha I} &= (\delta_{IJ} \delta_{\alpha\beta} + M[v, \partial, \gamma, \gamma_5; \theta]_{\alpha\beta IJ}) \psi_{\beta J} \\ \bar{\Psi}_{\alpha I} &= (\delta_{IJ} \delta_{\alpha\beta} + \bar{M}[v, \partial, \gamma, \gamma_5; \theta]_{\alpha\beta IJ}) \bar{\psi}_{\beta J} \\ [d\bar{\psi}] [d\psi] &= \det (\mathbb{I} + \bar{M}) \det (\mathbb{I} + M) [d\bar{\Psi}] [d\Psi] , \end{aligned} \quad (3.3)$$

where the determinants are defined by their diagrammatic expansion in dimensional regularization in powers of θ . Now, in dimensional regularization we have

$$\det (\mathbb{I} + M) = \det (\mathbb{I} + \bar{M}) = 1 . \quad (3.4)$$

To see this, take *e. g.* the determinant $\det (\mathbb{I} + M)$ and write it as the partition function

$$\det (\mathbb{I} + M) = \int [d\bar{\psi}] [d\psi] e^{iS[M]}$$

of a fermion theory with classical action

$$S[M] = \int d^{2\omega}x \bar{\psi} (\mathbb{I} + M) \psi .$$

The propagator of such a theory is the identity and the interaction vertices come from the operator $M[v, \partial, \gamma, \gamma_5; \theta]$, so the Feynman integrals that enter the diagrammatic expansion of $\det (\mathbb{I} + M)$ are of the form

$$\int \frac{d^{2\omega}q}{(2\pi)^{2\omega}} q_{\mu_1} \cdots q_{\mu_n} .$$

Since this integral vanishes in dimensional regularization, eq. (3.4) holds and the change of variables (3.3) gives for the path integral in (3.2)

$$Z^{\text{reg}}[v; \theta] = \mathcal{N} \int [d\bar{\Psi}] [d\Psi] \exp \left(i S_{\text{nc}}^{\text{reg}}[\bar{\Psi}, \Psi, V_\mu] \right) . \quad (3.5)$$

Hence $Z^{\text{reg}}[v; \theta]$ is a functional of V_μ , and so is the regularized effective action

$$\Gamma^{\text{reg}}[v; \theta] = -i \ln Z[v; \theta]^{\text{reg}} = \Gamma^{\text{reg}}[V] . \quad (3.6)$$

In other words, the regularized effective action depends on v_μ through V_μ .

Eq. (3.6) for $\Gamma^{\text{reg}}[V]$ is to be understood in a diagrammatic sense as the generating functional of 1PI Green functions for the field V_μ . That is to say,

$$i\Gamma^{\text{reg}}[V] = \sum_{n=1}^{\infty} \frac{1}{n!} \int d^{2\omega}x_1 \dots \int d^{2\omega}x_n V_{\mu_1 I_1 J_1}(x_1) \dots V_{\mu_n I_n J_n}(x_n) \Gamma_{I_1 J_1 \dots I_n J_n}^{\mu_1 \dots \mu_n}(x_1, \dots, x_n) , \quad (3.7)$$

with

$$\Gamma_{I_1 J_1 \dots I_n J_n}^{\mu_1 \dots \mu_n}(x_1, \dots, x_n) = \langle \mathcal{J}_{I_1 J_1}^{\mu_1}(x_1) \dots \mathcal{J}_{I_n J_n}^{\mu_n}(x_n) \rangle_0^{\text{conn}} \quad (3.8)$$

and

$$\mathcal{J}_{I_i J_i}^{\mu_i}(x_i) = (\Psi_{\beta_i J_i} \star \bar{\Psi}_{\alpha_i I_i})(x_i) (\bar{\gamma}^{\mu_i} \gamma_5)_{\alpha_i \beta_i} . \quad (3.9)$$

Here the symbol $\langle \mathcal{O} \rangle_0^{\text{conn}}$ stands for the connected component of the correlation function $\langle \mathcal{O} \rangle_0$ defined by

$$\langle \mathcal{O} \rangle_0 = \int [d\bar{\Psi}] [d\Psi] \mathcal{O} \exp \left(i \int d^{2\omega}x \bar{\Psi}_I \not{\partial} \Psi_I \right) . \quad (3.10)$$

Note that eqs. (3.8) and (3.10) define $\Gamma_{I_1 J_1 \dots I_n J_n}^{\mu_1 \dots \mu_n}(x_1, \dots, x_n)$ as the result of applying Wick's theorem to $\mathcal{J}_{I_1 J_1}^{\mu_1}(x_1) \dots \mathcal{J}_{I_n J_n}^{\mu_n}(x_n)$ with regard to the contraction

$$\langle \Psi_{\beta J}(y) \bar{\Psi}_{\alpha I}(x) \rangle_0 = \delta_{JI} \int \frac{d^{2\omega}q}{(2\pi)^{2\omega}} e^{-iq(y-x)} \frac{i \not{q}_{\beta\alpha}}{q^2 + i0^+} . \quad (3.11)$$

It is not difficult to see that in eq. (3.8) there are $(n-1)!$ different contractions and that, upon combination with the V 's in eq. (3.7), they all yield the same contribution. The regularized effective action then takes the form

$$i\Gamma^{\text{reg}}[V] = - \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \int d^{2\omega}x_1 \dots \int d^{2\omega}x_n \text{Tr} [V_{\mu_1}(x_1) \dots V_{\mu_n}(x_n)] \Gamma^{\mu_1 \dots \mu_n}(x_1, \dots, x_n) , \quad (3.12)$$

where

$$\text{Tr} [V_{\mu_1}(x_1) \dots V_{\mu_n}(x_n)] = V_{\mu_1 I_1 I_2}(x_1) V_{\mu_2 I_2 I_3}(x_2) \dots V_{\mu_{n-1} I_{n-1} I_n}(x_{n-1}) V_{\mu_n I_n I_1}(x_n) ,$$

the 1PI Green function $\Gamma_{\mu_1 \dots \mu_n}(x_1, \dots, x_n)$ reads

$$\begin{aligned} \Gamma_{\mu_1 \dots \mu_n}(x_1, \dots, x_n) &= i^n \int \prod_{i=1}^n \frac{d^{2\omega}p_i}{(2\pi)^{2\omega}} (2\pi)^{2\omega} \delta(p_1 + \dots + p_n) e^{i \sum_{i=1}^n p_i x_i} e^{-\frac{i}{2} \sum_{1 \leq i < j < n} \theta^{\alpha\beta} p_{i\alpha} p_{j\beta}} \\ &\times \int \frac{d^{2\omega}q}{(2\pi)^{2\omega}} \frac{\text{tr} [(\not{q} + \not{p}_1) \bar{\gamma}^{\mu_1} \text{P}_- \not{q} \bar{\gamma}^{\mu_2} \text{P}_- (\not{q} - \not{p}_2) \dots (\not{q} - \sum_{i=2}^{n-1} \not{p}_i) \bar{\gamma}^{\mu_n} \text{P}_-]}{(q + p_1)^2 q^2 (q - p_2)^2 \dots (q - \sum_{i=2}^{n-1} p_i)^2} \end{aligned} \quad (3.13)$$

and the symbol tr denotes trace over Dirac matrices. For completeness we present very briefly an alternative derivation of (3.12). Integrating over $[d\bar{\Psi}]$ and $[d\Psi]$ in eq. (3.5) and using eq. (3.6), we obtain

$$i\Gamma^{\text{reg}}[V] = \text{Tr} \ln [1 + (\not{\partial})^{-1} \bar{\gamma}^\mu V_\mu \text{P}_- \star] = - \sum_{n=1}^n \frac{(-1)^n}{n} \text{Tr} [(\not{\partial})^{-1} \bar{\gamma}^\mu V_\mu \text{P}_- \star]^n, \quad (3.14)$$

where Tr is to be interpreted as $\int d^{2\omega}x$ for the continuous indices of the operator on which Tr acts and $(\not{\partial})^{-1}$ has matrix elements $\langle y | (\not{\partial})^{-1} | x \rangle$ given by the right-hand-side of eq. (3.11). Clearly the right-hand side of eq. (3.14) has a neat diagrammatic representation which readily leads to eq. (3.12).

We stress the fact that the noncommutative field $V_\mu(x)$ in eqs. (3.12) and (3.14) is a mere spectator in the sense that these equations hold whatever the algebra on which $V_\mu(x)$ takes values be, provided the operation $\mathbf{Tr}$ make sense. Eqs. (3.12) and (3.14) are thus valid for noncommutative $U(N)$, simple, semisimple and non-semisimple gauge groups. One then expects that for nonsemisimple gauge groups a renormalized effective action $\Gamma^{\text{ren}}[V]$ can be defined so that the noncommutative gauge anomaly has the same form as for noncommutative $U(N)$ group, *i.e.* such that

$$s_\star \Gamma^{\text{ren}}[V] = \mathcal{A}_\star, \quad (3.15)$$

with

$$\mathcal{A}_\star = -\frac{i}{24\pi^2} \int d^4x \epsilon^{\mu_1\mu_2\mu_3\mu_4} \mathbf{Tr} \Lambda \star \partial_{\mu_1} \left(V_{\mu_2} \star \partial_{\mu_3} V_{\mu_4} + \frac{1}{2} V_{\mu_2} \star V_{\mu_3} \star V_{\mu_4} \right). \quad (3.16)$$

In the remainder of the section we prove that is indeed so.

To demonstrate eqs. (3.15) and (3.16) we proceed as follows. Since the integral over $d^{2\omega}q$ in eq. (3.12) does not involve any nonplanar factor $e^{iq_\alpha \theta^{\alpha\beta} p_{i\beta}}$, the effective action in eq. (3.12) is given by a sum over dimensionally regularized planar diagrams. Hence, the Quantum action principle [22] holds for this effective action and the following equation is valid

$$s_\star \Gamma^{\text{reg}}[V] = \hat{\Delta} \cdot \Gamma^{\text{reg}}[V]. \quad (3.17)$$

Here $\hat{\Delta} \cdot \Gamma^{\text{reg}}[V]$ is the insertion in $\Gamma^{\text{reg}}[V]$ of the evanescent operator $\hat{\Delta}$ defined by

$$\hat{\Delta} = s_\star S_{\text{nc}}^{\text{reg}} = \int d^{2\omega}x \left[\bar{\Psi}_I \star \Lambda_{IJ} \star i \hat{\not{\partial}} \text{P}_+ \Psi_J - \bar{\Psi}_I \star i \hat{\not{\partial}} (\Lambda_{IJ} \star \text{P}_- \Psi_J) \right].$$

Substituting this in eq. (3.17), we obtain for its right-hand side

$$\begin{aligned} \hat{\Delta} \cdot \Gamma^{\text{reg}}[V] = & - \sum_{n=1}^{\infty} (-1)^n \int d^{2\omega} x \int d^{2\omega} x_1 \dots \int d^{2\omega} x_n \\ & \times \text{Tr} [\Lambda(x) V_{\mu_1}(x_1) \dots V_{\mu_n}(x_n)] \Gamma^{\mu_1 \dots \mu_n}(x, x_1, \dots, x_n \mid \hat{\Delta}) , \end{aligned} \quad (3.18)$$

where

$$\begin{aligned} \Gamma^{\mu_1 \dots \mu_n}(x, x_1, \dots, x_n \mid \hat{\Delta}) = & i^{n+1} \int \frac{d^{2\omega} p}{(2\pi)^{2\omega}} \int \prod_{i=1}^n \frac{d^{2\omega} p_i}{(2\pi)^{2\omega}} e^{i(p x + \sum_{i=1}^n p_i x_i)} \\ & \times e^{-\frac{i}{2} \sum_{1 \leq i < j \leq n} \theta^{\alpha\beta} p_{i\alpha} p_{j\beta}} (2\pi)^{2\omega} \delta(p + p_1 + \dots + p_n) \Gamma^{\mu_1 \dots \mu_n}(p, p_1, \dots, p_n \mid \hat{\Delta}) , \end{aligned} \quad (3.19)$$

and the 1PI Green function $\Gamma^{\mu_1 \dots \mu_n}(p, p_1, \dots, p_n \mid \hat{\Delta})$ with the insertion reads

$$\begin{aligned} \Gamma^{\mu_1 \dots \mu_n}(p, p_1, \dots, p_n \mid \hat{\Delta}) = & \int \frac{d^{2\omega} q}{(2\pi)^{2\omega}} \frac{1}{q^2 (q - p_1)^2 (q - p_1 - p_2)^2 \dots (q - \sum_{i=1}^n p_i)^2} \\ & \times \text{tr} [\hat{\not{q}} P_+ - (\hat{\not{q}} - \sum_{i=1}^n \hat{\not{p}}_i) P_-] \hat{\not{q}} \bar{\gamma}^{\mu_1} P_- (\hat{\not{q}} - \hat{\not{p}}_1) \bar{\gamma}^{\mu_2} P_- (\hat{\not{q}} - \hat{\not{p}}_1 - \hat{\not{p}}_2) \dots \bar{\gamma}^{\mu_n} P_- (\hat{\not{q}} - \sum_{i=1}^n \hat{\not{p}}_i) . \end{aligned} \quad (3.20)$$

As before, tr denotes trace over Dirac matrices. For $n \geq 5$ the integral in eq. (3.20) is UV finite by power counting at $2\omega = 4$. Hence,

$$\Gamma^{\mu_1 \dots \mu_n}(p, p_1, \dots, p_n \mid \hat{\Delta}) = O(\varepsilon) \quad n \geq 5 , \quad (3.21)$$

where $\varepsilon = \omega - 2$. As concerns $n \leq 4$, using the results in Appendix A, it is straightforward to compute the contribution $\Gamma_{\text{eps}}^{\mu_1 \dots \mu_n}(p, p_1, \dots, p_n \mid \hat{\Delta})$ to $\Gamma^{\mu_1 \dots \mu_n}(p, p_1, \dots, p_n \mid \hat{\Delta})$ involving $\epsilon^{\mu_1 \mu_2 \mu_3 \mu_4}$. After some calculations, we obtain

$$\begin{aligned} \Gamma_{\text{eps}}^{\mu_1}(p, p_1 \mid \hat{\Delta}) &= 0 \\ \Gamma_{\text{eps}}^{\mu_1 \mu_2}(p, p_1, p_2 \mid \hat{\Delta}) &= \frac{1}{24\pi^2} \epsilon^{\rho\mu_1\sigma\mu_2} p_{1\rho} p_{2\sigma} + O(\varepsilon) \\ \Gamma_{\text{eps}}^{\mu_1 \mu_2 \mu_3}(p, p_1, p_2, p_3 \mid \hat{\Delta}) &= -\frac{1}{2} \frac{1}{24\pi^2} \epsilon^{\rho\mu_1\mu_2\mu_3} (p_1 + p_2 + p_3)_\rho + O(\varepsilon) \\ \Gamma_{\text{eps}}^{\mu_1 \mu_2 \mu_3 \mu_4}(p, p_1, p_2, p_3, p_4 \mid \hat{\Delta}) &= O(\varepsilon) . \end{aligned} \quad (3.22)$$

Substituting eqs. (3.21) and (3.22) in eq. (3.19), and the result so obtained in eq. (3.18), we have that the contribution to the right-hand side of eq. (3.17) which contains $\epsilon^{\mu_1 \mu_2 \mu_3 \mu_4}$ reads

$$\hat{\Delta} \cdot \Gamma^{\text{reg}}[V] \Big|_{\text{eps}} = \mathcal{A}_\star^{\text{reg}} , \quad (3.23)$$

where

$$\mathcal{A}_\star^{\text{reg}} = -\frac{i}{24\pi^2} \int d^2\omega x \epsilon^{\mu_1\mu_2\mu_3\mu_4} \text{Tr} \Lambda \star \partial_{\mu_1} \left(V_{\mu_2} \star \partial_{\mu_3} V_{\mu_4} + \frac{1}{2} V_{\mu_2} \star V_{\mu_3} \star V_{\mu_4} \right) + O(\varepsilon) .$$

Hence, if $\Gamma_{\text{eps}}^{\text{reg}}[V]$ denotes the contribution to the regularized effective action $\Gamma^{\text{reg}}[V]$ involving $\epsilon^{\mu_1\mu_2\mu_3\mu_4}$, eqs. (3.17) and (3.23) imply

$$s_\star \Gamma_{\text{eps}}^{\text{reg}}[V] = \hat{\Delta} \cdot \Gamma^{\text{reg}}[V] \Big|_{\text{eps}} = \mathcal{A}_\star^{\text{reg}} . \quad (3.24)$$

It is not difficult to show that the pole part of $\Gamma^{\text{reg}}[V]$ at $\varepsilon = 0$ does not depend on $\epsilon^{\mu_1\mu_2\mu_3\mu_4}$. This, together with the observation that any vector-like contribution to the regularized effective action – *i.e* not involving $\epsilon^{\mu_1\mu_2\mu_3\mu_4}$ – can be regularized in a gauge invariant way within the framework of dimensional regularization, implies that it is always possible to define a renormalized effective action

$$\Gamma^{\text{ren}}[V] = \Gamma_{\text{vec-like}}^{\text{ren}}[V] + \Gamma_{\text{eps}}^{\text{ren}}[V]$$

such that

$$s_\star \Gamma_{\text{vec-like}}^{\text{ren}}[V] = 0$$

and

$$s_\star \Gamma_{\text{eps}}^{\text{ren}}[V] = \lim_{\varepsilon \rightarrow 0} \mathcal{A}_\star^{\text{reg}} = \mathcal{A}_\star ,$$

with $\mathcal{A}_\star$ as in eq. (3.16). Hence eqs. (3.15) and (3.16) follow.

Using finally that $s_\star V_\mu = sV_\mu$ and that V_μ is a function of v_μ and $\theta^{\mu\nu}$ we conclude that

$$s\Gamma^{\text{ren}}[v; \theta] = \mathcal{A}_\star . \quad (3.25)$$

This equation gives a simple expression for the anomaly if written in terms of the noncommutative fields V_μ and λ . In fact, $\mathcal{A}_{\text{nc}}^{\text{Bardeen}}$ in eq. (3.16) is nothing but the noncommutative counterpart of Bardeen's ordinary anomaly. However, in terms of the fields v_μ and λ , the anomaly is a complicated power series in $\theta^{\mu\nu}$ with coefficients depending on such fields. The first term of such series is the standard Bardeen anomaly $\mathcal{A}^{\text{Bardeen}}$ of ordinary spacetime,

$$\mathcal{A}_\star = \mathcal{A}^{\text{Bardeen}} + O(\theta) \quad (3.26)$$

$$\mathcal{A}^{\text{Bardeen}} = \mathcal{A}_\star|_{\theta=0} = -\frac{i}{24\pi^2} \int d^4x \epsilon^{\mu_1\mu_2\mu_3\mu_4} \text{Tr} \lambda \partial_{\mu_1} \left(v_{\mu_2} \partial_{\mu_3} v_{\mu_4} + \frac{1}{2} v_{\mu_2} v_{\mu_3} v_{\mu_4} \right) . \quad (3.27)$$

4 BRS triviality of θ -dependent contributions

The functional $\mathcal{A}_\star$ in eqs. (3.15) and (3.16) has been found by explicitly computing to all orders in $\theta^{\mu\nu}$ the $\epsilon^{\mu_1\mu_2\mu_3\mu_4}$ part of the one-loop radiative corrections to all the 1PI Green functions of the field V_μ . As is well known, only radiative corrections which are cohomologically nontrivial with respect to the ordinary chiral BRS operator s , that is to say, that can not be written as the s of something, yield a true anomalous contribution. To find the true anomaly, we must therefore identify in $\mathcal{A}_\star$ the cohomologically nontrivial contributions with respect to s . This we do next.

If in sections 2 and 3 we take as noncommutativity matrix $t\theta^{\mu\nu}$, with t a real parameter, we end up with a noncommutative BRS chiral operator $s_\star^{(t\theta)}$ and an anomaly $\mathcal{A}_\star^{(t\theta)}$ whose expressions are obtained from those in sections 2 and 3 by replacing $\theta^{\mu\nu}$ with $t\theta^{\mu\nu}$. Note that the dependence on t of $s_\star^{(t\theta)}$ is only through the Moyal product, which now is with respect to $t\theta^{\mu\nu}$, but that no explicit t -dependence is introduced (see ref. [21]). In Appendix B we prove that the logarithmic differential with respect to t of $\mathcal{A}_\star^{(t\theta)}$ is $s_\star^{(t\theta)}$ trivial, or in other words, that there exists a functional $\mathcal{B}[V^{(t\theta)}, t\theta]$ such that

$$t \frac{d}{dt} \mathcal{A}_\star^{(t\theta)} = s_\star^{(t\theta)} \mathcal{B}[V^{(t\theta)}, t\theta] . \quad (4.1)$$

Let us remark that we use the logarithmic derivative $t \frac{d}{dt}$, and not the ordinary derivative $\frac{d}{dt}$ as in refs. [23, 21], to be able to write everything in terms of the noncommutativity matrix $t\theta^{\mu\nu}$ and to avoid having to use both $\theta^{\mu\nu}$ and $t\theta^{\mu\nu}$. Integrating eq. (4.1) over t from 0 to 1 and using that –by definition of the Seiberg-Witten map– $s_\star^{(t\theta)} V_\mu^{(t\theta)} = s V_\mu^{(t\theta)}[v, t\theta]$, we have

$$\int_0^1 dt \frac{d\mathcal{A}_\star^{(t\theta)}}{dt} = \int_0^1 \frac{dt}{t} s_\star^{(t\theta)} \mathcal{B}[V^{(t\theta)}, t\theta] = \int_0^1 \frac{dt}{t} s \mathcal{B}[V^{(t\theta)}[v, t\theta], t\theta] .$$

Recalling now that $\mathcal{A}_\star^{(t\theta)} = \mathcal{A}_\star$ if $t = 1$ and $\mathcal{A}_\star^{(t\theta)} = \mathcal{A}^{\text{Bardeen}}$ if $t = 0$, and noting that the ordinary BRS chiral operator s does not depend on t , we obtain

$$\mathcal{A}_\star = \mathcal{A}^{\text{Bardeen}} - s \int_0^1 \frac{dt}{t} \mathcal{B}[V^{(t\theta)}[v, t\theta], t\theta] . \quad (4.2)$$

Hence the functional $\mathcal{A}_\star$ found in section 3 consists of two contributions: the standard Bardeen anomaly $\mathcal{A}^{\text{Bardeen}}$ of commutative spacetime, and a contribution –given by the second term in eq. (4.2)– which is cohomologically trivial with respect to the ordinary chiral BRS operator s . Comparing with eq. (3.26), we conclude that all contributions to $\mathcal{A}_\star$ of order one or higher

in $\theta^{\mu\nu}$ are cohomologically trivial, hence harmless, since they can be absorbed by adding finite counterterms to the renormalized effective action. Indeed, consider a new renormalized effective action $\Gamma'^{\text{ren}}[v; \theta]$ defined by

$$\Gamma'^{\text{ren}}[v; \theta] = \Gamma^{\text{ren}}[v; \theta] - \int_0^1 \frac{dt}{t} \mathcal{B}[V^{(t\theta)}[v, t\theta], t\theta] . \quad (4.3)$$

According to our discussion above, it follows that

$$s \Gamma'^{\text{ren}}[v; \theta] = \mathcal{A}^{\text{Bardeen}} .$$

We thus conclude that the anomaly is $\theta^{\mu\nu}$ -independent and has Bardeen's form.

5 Conclusion

In this paper we have calculated the chiral one-loop anomaly in 4-dimensional noncommutative gauge theories with arbitrary compact gauge group defined through the Seiberg-Witten map. Our main result is that for all these theories the chiral anomaly is the same as for their commutative counterparts. Hence any noncommutative chiral gauge theory of this type is anomaly free to one-loop order if, and only if, its ordinary counterpart is. This implies in particular that the anomaly cancellation conditions for the noncommutative standard model [6] and the noncommutative $SU(5)$ and $SO(10)$ models [13] are the same as for the ordinary ones [27]. We would like to emphasize that we have not found anomaly candidates but actually computed the anomaly, since we have calculated the relevant Feynman diagrams that produce the anomaly.

There is one key ingredient in our proof, namely that counterterms with mass dimension greater than four should be allowed in the renormalized effective action. This is necessary to cancel radiative corrections which, on the one hand, do not satisfy the equation $s\Gamma^{\text{ren}} = 0$ but, on the other, are cohomologically trivial with respect to s . This indicates that the proper framework for these theories is the effective field theory formalism, a proposal that has already been made by a number of authors [13, 17, 18]. If one insists on power-counting renormalizability, then the “safe” representations and the safe “groups” of ordinary gauge theories [24] are totally unsafe for noncommutative gauge theories, since they lead to anomalous theories [18].

Acknowledgments

CPM and FRR are grateful to CICyT, Spain for partial support through grant No. BFM2002-00950.

A Appendix: Useful integrals

To obtain the $\epsilon^{\mu_1\mu_2\mu_3\mu_4}$ contribution to the n -point functions $\Gamma^{\mu_1\cdots\mu_n}(p, p_1, \dots, p_n \mid \hat{\Delta})$ with one evanescent insertion $\hat{\Delta}$ given in eqs. (3.22) the following integrals are needed:

$$\begin{aligned}
\int \frac{d^{2\omega}q}{(2\pi)^{2\omega}} \frac{\hat{q}^2}{q^2 (q - q_1)^2 (q - p_2)^2} &= -\frac{1}{2} \frac{i}{16\pi^2} + O(\varepsilon) \\
\int \frac{d^{2\omega}q}{(2\pi)^{2\omega}} \frac{\hat{q}^2 \bar{q}_\mu}{q^2 (q - q_1)^2 (q - q_2)^2} &= -\frac{1}{6} \frac{i}{16\pi^2} (\bar{q}_1 + \bar{q}_2)_\mu + O(\varepsilon) \\
\int \frac{d^{2\omega}q}{(2\pi)^{2\omega}} \frac{\hat{q}^2 \bar{q}_{\mu_1} \bar{q}_{\mu_2}}{q^2 (q - q_1)^2 (q - q_2)^2 (q - q_3)^2} &= -\frac{1}{12} \frac{i}{16\pi^2} \bar{g}_{\mu_1\mu_2} + O(\varepsilon) \\
\int \frac{d^{2\omega}q}{(2\pi)^{2\omega}} \frac{\hat{q}^2 \bar{q}_{\mu_1} \bar{q}_{\mu_2} \bar{q}_{\mu_3}}{q^2 (q - q_1)^2 (q - q_2)^2 (q - q_3)^2} &= \\
&= -\frac{1}{48} \frac{i}{16\pi^2} \sum_{i=1}^3 (\bar{g}_{\mu_1\mu_2} \bar{q}_{i\mu_3} + \bar{g}_{\mu_1\mu_3} \bar{q}_{i\mu_2} + \bar{g}_{\mu_2\mu_3} \bar{q}_{i\mu_1}) + O(\varepsilon) \\
\int \frac{d^{2\omega}q}{(2\pi)^{2\omega}} \frac{\hat{q}^2 \bar{q}^2 \bar{q}_{\mu_1} \bar{q}_{\mu_2}}{q^2 (q - q_1)^2 (q - q_2)^2 (q - q_3)^2 (q - q_4)^2} &= -\frac{1}{16} \frac{i}{16\pi^2} \bar{g}_{\mu_1\mu_2} + O(\varepsilon) .
\end{aligned}$$

Here $\varepsilon = \omega - 2$.

B Appendix: Proof of eq. (4.1)

In what follows we will use $\omega^{\alpha\beta}$ for $t\theta^{\alpha\beta}$, denote the Moyal product with respect to $\omega^{\alpha\beta}$ by $\star$ and write a small circle $\circ$ for the logarithmic differential with respect to t , *i.e.*

$$\omega^{\alpha\beta} = t\theta^{\alpha\beta} \quad \star = \star_\omega \quad \mathcal{F}^\circ = t \frac{d\mathcal{F}}{dt} . \quad (\text{B.1})$$

The functional $\mathcal{A}_\star^{(t\theta)}$, which in this notation we write as $\mathcal{A}_\star$, has a piece of order zero in $\omega^{\alpha\beta}$ given by $\mathcal{A}^{\text{Bardeen}}$ in eq. (3.27) and a piece that collects all the higher order terms in $\omega^{\alpha\beta}$ and

which precisely gives the contributions to $\overset{\circ}{\mathcal{A}}_\star$. We want to prove eq. (4.1), which now takes the form

$$\overset{\circ}{\mathcal{A}}_\star = s_\star \mathcal{B} . \quad (\text{B.2})$$

Using

$$f \overset{\circ}{\star} g = \frac{1}{2} \omega^{\alpha\beta} \partial_\alpha f \star \partial_\beta g \quad (\text{B.3})$$

and [23]

$$\overset{\circ}{V}_\mu = -\frac{i}{4} \omega^{\alpha\beta} \{V_\alpha, F_{\beta\mu} + \partial_\beta V_\mu\}_\star \quad \overset{\circ}{\Lambda} = \frac{i}{4} \omega^{\alpha\beta} \{\partial_\alpha \Lambda, V_\beta\}_\star , \quad (\text{B.4})$$

the functional $\overset{\circ}{\mathcal{A}}_\star$ can be expanded as a sum

$$\overset{\circ}{\mathcal{A}}_\star = \overset{\circ}{\mathcal{A}}_{\star,3} + \overset{\circ}{\mathcal{A}}_{\star,4} + \overset{\circ}{\mathcal{A}}_{\star,5} + \overset{\circ}{\mathcal{A}}_{\star,6} , \quad (\text{B.5})$$

where $\overset{\circ}{\mathcal{A}}_{\star,n}$ collects all contributions in $\overset{\circ}{\mathcal{A}}_\star$ of degree n in the fields Λ and V_μ (see below for their explicit expressions). In turn, the noncommutative chiral BRS operator $s_\star$ can be written as the sum

$$s_\star = s_{\star,0} + s_{\star,1} \quad (\text{B.6})$$

of two operators $s_{\star,0}$ and $s_{\star,1}$ whose action on the fields Λ and V_μ is given by

$$s_{\star,0} V_\mu = \partial_\mu \Lambda \quad s_{\star,0} \Lambda = 0 \quad (\text{B.7})$$

$$s_{\star,1} V_\mu = [V_\mu, \Lambda]_\star \quad s_{\star,1} \Lambda = -\Lambda \star \Lambda . \quad (\text{B.8})$$

These two operators satisfy

$$s_{\star,0}^2 = 0 \quad s_{\star,0} s_{\star,1} + s_{\star,1} s_{\star,0} = 0$$

and have the important property that $s_{\star,0}$ preserves the degree in the fields and $s_{\star,1}$ increases it by one. From eqs. (B.5) and (B.6) it follows that to prove eq. (B.2) it is sufficient to take for $\mathcal{B}$ an expansion

$$\mathcal{B} = \mathcal{B}_3 + \mathcal{B}_4 + \mathcal{B}_5 + \mathcal{B}_6$$

in the number of fields and show that

$$\overset{\circ}{\mathcal{A}}_{\star,3} = s_{\star,0} \mathcal{B}_3 \quad (\text{B.9})$$

$$\overset{\circ}{\mathcal{A}}_{\star,4} - s_{\star,1} \mathcal{B}_3 = s_{\star,0} \mathcal{B}_4 \quad (\text{B.10})$$

$$\overset{\circ}{\mathcal{A}}_{\star,5} - s_{\star,1} \mathcal{B}_4 = s_{\star,0} \mathcal{B}_5 \quad (\text{B.11})$$

$$\overset{\circ}{\mathcal{A}}_{\star,6} - s_{\star,1} \mathcal{B}_5 = s_{\star,0} \mathcal{B}_6 \quad (\text{B.12})$$

$$s_{\star,1} \mathcal{B}_6 = 0 . \quad (\text{B.13})$$

Hence, to prove (B.2) all we have to do is finding functionals $\mathcal{B}_3, \mathcal{B}_4, \mathcal{B}_5$ and $\mathcal{B}_6$ satisfying the ladder equations. To do this it is convenient to use differential forms, so let us write eqs. (B.7) and (B.8) in terms of differential forms. Recalling that $V = V_\mu dx^\mu$ and using $\{dx^\mu, s_{\star,0}\} = \{dx^\mu, s_{\star,1}\} = \{\Lambda, dx^\mu\} = 0$, we have

$$s_{\star,0}V = -d\Lambda \quad s_{\star,0}\Lambda = 0 \quad (\text{B.14})$$

$$s_{\star,1}V = -\{V, \Lambda\}_\star \quad s_{\star,1}\Lambda = -\Lambda \star \Lambda . \quad (\text{B.15})$$

B.1 Computation of $\mathcal{B}_3$ and $\mathcal{B}_4$

Taking the logarithmic differential with respect to t of $\mathcal{A}_\star$ and using eqs. (B.3) and (B.4), it is straightforward to see that

$$\mathring{\mathcal{A}}_{\star,3} = -\frac{i}{24\pi^2} \int \frac{i}{2} \omega^{\alpha\beta} \text{Tr} \partial_\alpha \Lambda \star \partial_\beta dV \star dV .$$

It is clear that

$$\mathcal{B}_3 = -\frac{i}{24\pi^2} \int \frac{i}{2} \omega^{\alpha\beta} \text{Tr} [x V_\alpha \star \partial_\beta dV \star dV - (1-x) V_\alpha \star dV \star \partial_\beta dV] , \quad (\text{B.16})$$

with x an arbitrary parameter, solves eq. (B.9). Indeed, acting with $s_{\star,0}$ on $\mathcal{B}_3$, integrating by parts the derivative ∂_β in the second term in eq. (B.16) and neglecting the integral of a divergence, we recover $\mathring{\mathcal{A}}_{\star,3}$. Note that eq. (B.16) provides a one-parameter family of solutions for $\mathcal{B}_3$. Furthermore, to $\mathcal{B}_3$ we can also add a term

$$\int \omega^{\alpha\beta} \text{Tr} \partial_\alpha V_\beta \star dV \star dV$$

with arbitrary coefficient, since $s_{\star,0}$ acting on it vanishes.

Let us move now on to eq. (B.10). We first calculate $\mathring{\mathcal{A}}_{\star,4}$ and $s_{\star,1}\mathcal{B}_3$. Acting with $t \frac{d}{dt}$ on $\mathcal{A}_\star$, noting eqs. (B.3) and (B.4), retaining terms of order four in the fields, using the cyclic property of the trace Tr and of the integral of a Moyal product of functions to push the ghost field Λ to the far left, and integrating by parts whatever partial and/or exterior derivatives

act on Λ , we obtain after some lengthy algebra that

$$\begin{aligned}\mathring{\mathcal{A}}_{\star,4} = & -\frac{i}{24\pi^2} \int \frac{i}{4} \omega^{\alpha\beta} \mathbf{Tr} \Lambda \star \left[V_\alpha \star dV \star \partial_\beta dV - V_\alpha \star \partial_\beta dV \star dV + V \star \partial_\alpha V \star \partial_\beta dV \right. \\ & - V \star \partial_\alpha dV \star \partial_\beta V - \partial_\alpha V_\beta \star dV \star dV - dV \star dV \star \partial_\alpha V_\beta - 2 \partial_\alpha V \star dV_\beta \star dV \\ & + 2 dV_\alpha \star dV_\beta \star dV - 2 dV_\alpha \star \partial_\beta V \star dV - 2 dV \star \partial_\alpha V \star dV_\beta + 2 dV \star dV_\alpha \star dV_\beta \\ & - 2 dV \star dV_\alpha \star \partial_\beta V + dV \star \partial_\alpha V \star \partial_\beta V - \partial_\alpha V \star \partial_\beta V \star dV - \partial_\alpha V \star dV \star \partial_\beta V \\ & - \partial_\alpha dV \star dV \star V_\beta + 2 \partial_\alpha dV \star V_\beta \star dV + dV \star \partial_\alpha dV \star V_\beta - 2 dV \star V_\alpha \star \partial_\beta dV \\ & \left. + \partial_\alpha dV \star \partial_\beta V \star V - \partial_\alpha V \star \partial_\beta dV \star V + \partial_\alpha dV \star V \star \partial_\beta V + \partial_\alpha V \star V \star \partial_\beta dV \right]\end{aligned}$$

Proceeding similarly for $s_{\star,1}\mathcal{B}_3$, and taking for simplicity $x = 1$, we have

$$\begin{aligned}s_{\star,1}\mathcal{B}_3 = & \frac{i}{24\pi^2} \int \frac{i}{2} \omega^{\alpha\beta} \mathbf{Tr} \Lambda \star \left[V_\alpha \star \partial_\beta dV \star dV + V \star dV_\alpha \star \partial_\beta dV \right. \\ & + V \star dV \star \partial_\alpha dV_\beta + V \star \partial_\alpha dV \star dV_\alpha + dV_\alpha \star \partial_\beta dV \star V \\ & \left. + dV \star \partial_\alpha dV_\beta \star V + \partial_\alpha dV \star dV \star V_\beta + \partial_\alpha dV \star dV_\beta \star V \right]\end{aligned}$$

To simplify these expressions we introduce the notation

$$A^i = -\frac{i}{24\pi^2} \int \frac{i}{4} \omega^{\alpha\beta} \mathbf{Tr} \Lambda \star a_{\alpha\beta}^i ,$$

with $a_{\alpha\beta}^i$ as in Table 1. Note that $\omega^{\alpha\beta} a_{\alpha\beta}^i$ is a 4-form with one explicit $\omega^{\alpha\beta}$, three explicit derivatives and three noncommutative gauge fields. By “explicit” here we mean ω ’s and ∂ ’s that are not hidden in the $\star$ -product. In Table 1 we have listed all such forms that can be constructed. With this notation $\mathring{\mathcal{A}}_{\star,4} - s_{\star,1}\mathcal{B}$ reads

$$\begin{aligned}\mathring{\mathcal{A}}_{\star,4} - s_{\star,1}\mathcal{B} = & A^1 - A^2 + A^3 + A^4 - A^5 + A^6 + A^7 - A^8 + A^9 - A^{10} - A^{12} \\ & - 2A^{13} - 2A^{15} - 2A^{16} - 2A^{18} + A^{19} + A^{20} - 2A^{21} + 2A^{22} + A^{23} \\ & + A^{24} + 2A^{26} + 2A^{27} + 2A^{28} + 2A^{29} + 2A^{32} + 2A^{36} + 2A^{37} + 2A^{39} .\end{aligned}\tag{B.17}$$

Now, not all the 4-forms $\omega^{\alpha\beta} a_{\alpha\beta}^i$ in Table 1 are linearly independent. To see this, consider *e.g.* the 5-form $\Omega^\beta = \omega^{\alpha\beta} \partial_\alpha V \star dV \star dV$ and act on it with the inner contraction

$$i_\beta \equiv i_{\partial_\beta} = \frac{\partial}{\partial(dx^\beta)} .$$

$a_{\alpha\beta}^1$	$dV \star \partial_\alpha V \star \partial_\beta V$	$a_{\alpha\beta}^{14}$	$\partial_\alpha V \star dV \star dV_\beta$	$a_{\alpha\beta}^{27}$	$V \star dV_\alpha \star \partial_\beta dV$
$a_{\alpha\beta}^2$	$\partial_\alpha V \star dV \star \partial_\beta V$	$a_{\alpha\beta}^{15}$	$\partial_\alpha V \star dV_\beta \star dV$	$a_{\alpha\beta}^{28}$	$\partial_\alpha dV \star dV_\beta \star V$
$a_{\alpha\beta}^3$	$\partial_\alpha V \star \partial_\beta V \star dV$	$a_{\alpha\beta}^{16}$	$dV \star dV_\alpha \star \partial_\beta V$	$a_{\alpha\beta}^{29}$	$dV_\alpha \star \partial_\beta dV \star V$
$a_{\alpha\beta}^4$	$\partial_\alpha dV \star V \star \partial_\beta V$	$a_{\alpha\beta}^{17}$	$dV_\alpha \star dV \star \partial_\beta V$	$a_{\alpha\beta}^{30}$	$dV_\alpha \star V \star \partial_\beta dV$
$a_{\alpha\beta}^5$	$V \star \partial_\alpha dV \star \partial_\beta V$	$a_{\alpha\beta}^{18}$	$dV_\alpha \star \partial_\beta V \star dV$	$a_{\alpha\beta}^{31}$	$\partial_\alpha dV_\beta \star dV \star V$
$a_{\alpha\beta}^6$	$V \star \partial_\alpha V \star \partial_\beta dV$	$a_{\alpha\beta}^{19}$	$\partial_\alpha dV \star dV \star V_\beta$	$a_{\alpha\beta}^{32}$	$dV \star \partial_\alpha dV_\beta \star V$
$a_{\alpha\beta}^7$	$\partial_\alpha dV \star \partial_\beta V \star V$	$a_{\alpha\beta}^{20}$	$dV \star \partial_\alpha dV \star V_\beta$	$a_{\alpha\beta}^{33}$	$dV \star V \star \partial_\alpha dV_\beta$
$a_{\alpha\beta}^8$	$\partial_\alpha V \star \partial_\beta dV \star V$	$a_{\alpha\beta}^{21}$	$dV \star V_\alpha \star \partial_\beta dV$	$a_{\alpha\beta}^{34}$	$\partial_\alpha dV_\beta \star V \star dV$
$a_{\alpha\beta}^9$	$\partial_\alpha V \star V \star \partial_\beta dV$	$a_{\alpha\beta}^{22}$	$\partial_\alpha dV \star V_\beta \star dV$	$a_{\alpha\beta}^{35}$	$V \star \partial_\alpha dV_\beta \star dV$
$a_{\alpha\beta}^{10}$	$dV \star dV \star \partial_\alpha V_\beta$	$a_{\alpha\beta}^{23}$	$V_\alpha \star \partial_\beta dV \star dV$	$a_{\alpha\beta}^{36}$	$V \star dV \star \partial_\alpha dV_\beta$
$a_{\alpha\beta}^{11}$	$dV \star \partial_\alpha V_\beta \star dV$	$a_{\alpha\beta}^{24}$	$V_\alpha \star dV \star \partial_\beta dV$	$a_{\alpha\beta}^{37}$	$dV \star dV_\alpha \star dV_\beta$
$a_{\alpha\beta}^{12}$	$\partial_\alpha V_\beta \star dV \star dV$	$a_{\alpha\beta}^{25}$	$\partial_\alpha dV \star V \star dV_\beta$	$a_{\alpha\beta}^{38}$	$dV_\alpha \star dV \star dV_\beta$
$a_{\alpha\beta}^{13}$	$dV \star \partial_\alpha V \star dV_\beta$	$a_{\alpha\beta}^{26}$	$V \star \partial_\alpha dV \star dV_\beta$	$a_{\alpha\beta}^{39}$	$dV_\alpha \star dV_\beta \star dV$

Table 1: All 4-forms with three derivatives and three gauge fields.

Being a 5-form in four dimensions, Ω^β is identically zero, and so is i_β acting on it. Hence

$$\begin{aligned}
0 &= i_\beta (\omega^{\alpha\beta} \partial_\alpha V \star dV \star dV) \\
&= \omega^{\alpha\beta} [\partial_\alpha V_\beta \star dV \star dV - \partial_\alpha V \star (\partial_\beta V - dV_\beta) \star dV - \partial_\alpha V \star dV \star (\partial_\beta V - dV_\beta)] \\
&= \omega^{\alpha\beta} (a_{\alpha\beta}^{12} - a_{\alpha\beta}^3 + a_{\alpha\beta}^{15} - a_{\alpha\beta}^2 + a_{\alpha\beta}^{14}) ,
\end{aligned}$$

which implies the relation

$$A^{12} - A^3 + A^{15} - A^2 + A^{14} = 0 .$$

This suggests that, to generate all the linear relations among the functionals A^i , it is enough to act with i_β on all the 5-forms Ω^β with one explicit $\omega^{\alpha\beta}$, three explicit derivatives and three noncommutative gauge fields. In listing the forms Ω^β , two restrictions should be observed. The first one is that it is only necessary to consider 5-forms Ω^β with at most two explicit

derivatives acting on the same field, since in Table 1 there is no $a_{\alpha\beta}^i$ with more than two explicit derivatives on the same gauge field. The second one is that whenever two explicit derivatives act on the same gauge field, they should not be both exterior derivatives. The reason for this is that 5-forms Ω^β containing an explicit d^2 do not provide, upon acting on them with i_β , any relation among the $\omega^{\alpha\beta}a_{\alpha\beta}^i$. Indeed, since $i_\beta d^2 = \partial_\beta d - d\partial_\beta$, the action of i_β on a 5-form containing an explicit d^2 yields a linear combination

$$i_\beta (5\text{-form with } d^2)^\beta = 4\text{-forms with } d^2 + 4\text{-form with } \partial_\beta d - d\partial_\beta$$

of 4-forms each of which is identically zero. There are twelve different forms Ω^α that can be constructed satisfying these restrictions on the derivatives, namely $\Omega^\beta = \omega^{\alpha\beta}\tilde{a}_\alpha$, with $\tilde{a}_\alpha$ given by

$$\begin{array}{cccc} \partial_\alpha V \star dV \star dV & dV_\alpha \star dV \star dV & \partial_\alpha dV \star dV \star V & \partial_\alpha dV \star V \star dV \\ dV \star \partial_\alpha V \star dV & dV \star dV_\alpha \star dV & dV \star \partial_\alpha dV \star V & V \star \partial_\alpha dV \star dV \\ dV \star dV \star \partial_\alpha V & dV \star dV \star dV_\alpha & dV \star V \star \partial_\alpha dV & V \star dV \star \partial_\alpha dV . \end{array}$$

If we act with i_β on these twelve 5-forms, we obtain the linear relations

$$\begin{array}{ll} A^2 + A^3 - A^{12} - A^{14} - A^{15} = 0 & A^7 + A^{19} - A^{28} - A^{31} = 0 \\ A^1 + A^3 - A^{11} - A^{13} - A^{18} = 0 & A^8 - A^{20} - A^{29} + A^{32} = 0 \\ A^1 + A^2 - A^{10} - A^{16} - A^{17} = 0 & A^9 + A^{21} - A^{30} - A^{33} = 0 \\ A^{12} + A^{17} + A^{18} - A^{38} - A^{39} = 0 & A^4 - A^{22} - A^{25} + A^{34} = 0 \\ A^{11} + A^{15} + A^{16} - A^{37} - A^{39} = 0 & A^5 + A^{23} - A^{26} - A^{35} = 0 \\ A^{10} + A^{13} + A^{14} - A^{37} - A^{38} = 0 & A^6 - A^{24} - A^{27} + A^{36} = 0 . \end{array}$$

Solving this system of equations for A^i ($i = 1, \dots, 12$) and substituting the solution in eq. (B.17), we write $\overset{\circ}{\mathcal{A}}_{\star,4} - s_{\star,1}\mathcal{B}_3$ in terms of the functionals A^i ($i = 13, \dots, 39$), the result being

$$\begin{aligned} \overset{\circ}{\mathcal{A}}_{\star,4} - s_{\star,1}\mathcal{B}_3 = & A^{13} + A^{14} - 4A^{15} - 4A^{16} + A^{17} + A^{18} - 3A^{21} + 3A^{22} \\ & + 2A^{23} + 2A^{24} + A^{25} + A^{26} + 3A^{27} + 3A^{28} + A^{29} + A^{30} \\ & + A^{31} + 3A^{32} + A^{33} - A^{34} - A^{35} + A^{36} + 2A^{37} - 3A^{38} + 2A^{39} . \end{aligned} \quad (\text{B.18})$$

We have thus obtained the left-hand side of eq. (B.10) in terms of linearly independent functionals A^i ($i = 13, \dots, 39$), each of which has one explicit $\omega^{\alpha\beta}$ and three explicit derivatives and has degree three in the noncommutative gauge field. It then follows that, for eq.

(B.10) to have a solution, $\mathcal{B}_4$ on the right-hand side must be a linear combination of functionals

$$B^r = -\frac{i}{24\pi^2} \int \frac{i}{4} \omega^{\alpha\beta} \mathbf{Tr} b_{\alpha\beta}^r, \quad (\text{B.19})$$

with $b_{\alpha\beta}^r$ a 4-form of order two in explicit derivatives and four in the noncommutative gauge field. With some patience, it can be seen that there are forty such functionals B^r whose $s_{\star,0}$ variation is not zero. Thirty of them can be written as linear combinations of the functionals

$b_{\alpha\beta}^1$	$V_\alpha \star V_\beta \star dV \star dV$	$b_{\alpha\beta}^6$	$\partial_\alpha V_\beta \star V \star dV \star V$
$b_{\alpha\beta}^2$	$dV_\alpha \star V_\beta \star dV \star V$	$b_{\alpha\beta}^7$	$\partial_\alpha V_\beta \star dV \star V \star V$
$b_{\alpha\beta}^3$	$dV \star dV_\alpha \star V \star V_\beta$	$b_{\alpha\beta}^8$	$V_\alpha \star dV \star \partial_\beta V \star V$
$b_{\alpha\beta}^4$	$dV_\alpha \star V_\beta \star V \star dV$	$b_{\alpha\beta}^9$	$V_\alpha \star \partial_\beta V \star V \star dV$
$b_{\alpha\beta}^5$	$\partial_\alpha V_\beta \star V \star V \star dV$	$b_{\alpha\beta}^{10}$	$V \star dV \star \partial_\alpha V \star V_\beta$

Table 2: All 4-forms with two derivatives and four gauge fields.

B^r whose $b_{\alpha\beta}^r$ are collected in Table 2. To illustrate that this is indeed so, let us consider as an example

$$B = -\frac{i}{24\pi^2} \int \frac{i}{4} \omega^{\alpha\beta} \mathbf{Tr} b_{\alpha\beta} \quad b_{\alpha\beta} = V \star V_\alpha \star \partial_\beta V \star dV.$$

Clearly, this $b_{\alpha\beta}$ is not in Table 2. However, using that

- (a) both $\mathbf{Tr}$ and the integral of a Moyal product of functions are cyclic,
- (b) that $\partial_\alpha = \{i_\alpha, d\}$, and
- (c) that $i_\beta(dV) \star dV \star V \star V_\alpha = -dV \star i_\beta(dV \star V \star V_\alpha)$,

and integrating by parts and neglecting total derivatives, we have

$$\begin{aligned}
B &\stackrel{(a,b)}{=} \frac{i}{24\pi^2} \int \frac{i}{4} \omega^{\alpha\beta} \mathbf{Tr} (i_\beta d + di_\beta) V \star dV \star V \star V_\alpha \\
&\stackrel{(c,d)}{=} -\frac{i}{24\pi^2} \int \frac{i}{4} \omega^{\alpha\beta} \mathbf{Tr} [dV \star i_\beta (dV \star V \star V_\alpha) + i_\beta V \star d(dV \star V \star V_\alpha)] \\
&\stackrel{(a)}{=} -\frac{i}{24\pi^2} \int \frac{i}{4} \omega^{\alpha\beta} \mathbf{Tr} (b_{\alpha\beta}^8 + b_{\alpha\beta}^3 + b_{\alpha\beta}^2) \\
&= B^8 + B^3 + B^2.
\end{aligned}$$

Similarly, any other functional B whose $b_{\alpha\beta}$ is not in Table 2 can be expressed as a linear combination of functionals B^r with $b_{\alpha\beta}^r$ in Table 2. It then follows that it is enough to write

for $\mathcal{B}_4$

$$\mathcal{B}_4 = \sum_{r=1}^{10} c_r B^r . \quad (\text{B.20})$$

To solve eq. (B.10) we need the $s_{\star,0}$ variation of $\mathcal{B}_4$. Acting with $s_{\star,0}$ on (B.20) and writing the result in terms of the linearly independent functionals A^i , corresponding to $i = 13, \dots, 39$, we obtain

$$\begin{aligned} s_{\star,0}\mathcal{B}_4 = & (-c_1 - c_2 + c_5 + c_6 + 2c_9) A^{13} \\ & + (-c_1 - c_4 + c_5 + c_6 + c_8 + 3c_9) A^{14} \\ & + (-2c_3 - c_5 + c_7 + c_8 + c_{10}) (A^{15} + A^{16}) \\ & + (c_1 - 2c_2 - c_4 - c_6 - c_7 + 2c_8 - c_9 + c_{10}) A^{17} \\ & + (c_1 - c_2 - 2c_4 - c_6 - c_7 + c_8 + c_{10}) A^{18} \\ & + (c_1 + c_8 - c_9) (A^{19} + A^{20}) \\ & + (c_3 - c_8) (-A^{21} + A^{22} + A^{27} + A^{28}) \\ & + (c_1 - c_2 - c_4) A^{23} \\ & + (c_1 - c_2 - c_4 + c_8 + c_{10}) A^{24} \\ & - c_2 (A^{25} + A^{29}) \\ & + (-c_4 + c_9 + c_{10}) (A^{26} + A^{30}) \\ & + (c_2 + c_7 - c_8) A^{31} \\ & + (c_3 + c_5) A^{32} \\ & + (c_2 - c_6 - c_9) A^{33} \\ & + (c_4 + c_6 + c_{10}) A^{34} \\ & + (c_3 - c_7 - c_8 - c_{10}) A^{35} \\ & + (c_4 - c_5 - c_9) A^{36} \\ & + (c_1 + c_3 - c_4 - c_6 - c_7 - c_9) A^{37} \\ & + (c_2 - c_3 + c_4 - c_5 + c_7 - c_9) A^{38} \\ & + (-c_1 + c_3 + c_4 + c_5 + c_6 + c_8 - c_9 + c_{10}) A^{39} . \end{aligned} \quad (\text{B.21})$$

Substituting now eqs. (B.18) and (B.21) in eq. (B.10) and equating the coefficients of A^i ($i = 13, \dots, 39$) on both sides, we obtain a system of 21 equations with unknowns $c_1, \dots, c_{10}$.

Its solution is

$$\begin{aligned} c_1 &= y + z & c_2 &= -1 & c_3 &= 3 - z & c_4 &= -1 + y + z & c_5 &= z \\ c_6 &= -y & c_7 &= 2 - z & c_8 &= -z & c_9 &= y & c_{10} &= z, \end{aligned}$$

where y and z are arbitrary parameters. This provides a two-parameter family of functionals $\mathcal{B}_4$ for which eq. (B.10) holds. Note that if we take $y = z = 0$, then $\mathcal{B}_4$ only has four terms.

B.2 Calculation of $\mathcal{B}_5$ and $\mathcal{B}_6$

One may proceed analogously as for $\mathcal{B}_4$ and explicitly compute $\mathcal{B}_5$ and $\mathcal{B}_6$. Here, instead, we present an alternative method which uses cohomological techniques. To apply them we shall employ the approach of ref. [23] which introduces gauge fields v_μ^A and ghost fields λ^A not only for the Lie algebra $\mathfrak{g}$ of the gauge group G but also for the whole enveloping algebra $\mathcal{U} = \{T_A\} = \{T_{\bar{a}}, T_i\}$ in which V_μ and Λ take values. Here the index $\bar{a}$ runs over the elements of $\mathfrak{g}$, so that in the notation of section 2 one has $\{T_{\bar{a}}\} = \{(T^k)^a, T^l\}$, while the index i runs over the complementary elements of $\mathcal{U}$. As shown in ref. [23], the standard Seiberg-Witten map can be extended to include $\mathcal{U}$ -valued fields v_μ and λ satisfying

$$sv_\mu^A = \partial_\mu \lambda^A + f_{BC}^A v_\mu^B \lambda^C \quad (\text{B.22})$$

$$s\lambda^A = \frac{1}{2} \lambda^B \lambda^C f_{CB}^A, \quad (\text{B.23})$$

with f_{AB}^C the structure constants of the Lie algebra $\mathcal{U}$, given by $[T_A, T_B] = f_{AB}^C T_C$. Of course, $\mathfrak{g}$ being a subalgebra of $\mathcal{U}$ means $f_{\bar{a}\bar{b}}^i = 0$ and implies that the BRS transformations above are subject to the truncation conditions

$$sv_\mu^A \Big|_{v_\mu^i = \lambda^i = 0} = \begin{cases} \partial_\mu \lambda^{\bar{a}} + f_{\bar{b}\bar{c}}^{\bar{a}} v_\mu^{\bar{b}} \lambda^{\bar{c}} & \text{if } A = \bar{a} \\ f_{\bar{b}\bar{c}}^i v_\mu^{\bar{b}} \lambda^{\bar{c}} = 0 & \text{if } A = i \end{cases} \quad (\text{B.24})$$

$$s\lambda^A \Big|_{v_\mu^i = \lambda^i = 0} = \begin{cases} \frac{1}{2} \lambda^{\bar{b}} \lambda^{\bar{c}} f_{\bar{b}\bar{c}}^{\bar{a}} & \text{if } A = \bar{a} \\ \frac{1}{2} \lambda^{\bar{b}} \lambda^{\bar{c}} f_{\bar{c}\bar{b}}^i = 0 & \text{if } A = i. \end{cases} \quad (\text{B.25})$$

The extended Seiberg-Witten map is defined by demanding

$$s_\star V_\mu^A = sV_\mu^A \quad s_\star \Lambda^A = s\Lambda^A, \quad (\text{B.26})$$

subject to the usual boundary conditions and with $s_\star$ defined by

$$s_\star V_\mu^A = \partial_\mu \Lambda^A + f_{BC}^A V_\mu^B \Lambda^C \quad s_\star \Lambda^A = \frac{1}{2} \Lambda^B \Lambda^C f_{CB}^A. \quad (\text{B.27})$$

By setting in it all fields v_μ^i and λ^i to zero, the standard Seiberg-Witten map is recovered. Furthermore, the truncation conditions imply that all formulas that hold for $\mathcal{U}$ -valued fields v_μ^A and λ^A will also hold for $\mathfrak{g}$ -valued fields $v_\mu^{\bar{a}}$ and $\lambda^{\bar{a}}$, and in particular eq. (B.2) that we want to prove. The idea is then to demonstrate eq. (B.2) for the extended Seiberg-Witten map.

We start from the fact that $\mathcal{A}_\star$ satisfies the anomaly consistency condition $s_\star \mathcal{A}_\star = 0$ which follows from eq. (B.2) because of $s_\star^2 = 0$. In terms of the commutative fields v_μ^A and λ^A , one has $s \mathcal{A}_\star = 0$. This implies $s \mathring{\mathcal{A}}_\star = 0$ since s commutes with the logarithmic derivative with respect to t . Using (B.26) again, one concludes $s_\star \mathring{\mathcal{A}}_\star = 0$ which decomposes into

$$s_{\star,0} \mathring{\mathcal{A}}_{\star,3} = 0 \quad (\text{B.28})$$

$$s_{\star,0} \mathring{\mathcal{A}}_{\star,4} + s_{\star,1} \mathring{\mathcal{A}}_{\star,3} = 0 \quad (\text{B.29})$$

$$s_{\star,0} \mathring{\mathcal{A}}_{\star,5} + s_{\star,1} \mathring{\mathcal{A}}_{\star,4} = 0 \quad (\text{B.30})$$

$$s_{\star,0} \mathring{\mathcal{A}}_{\star,6} + s_{\star,1} \mathring{\mathcal{A}}_{\star,5} = 0 \quad (\text{B.31})$$

$$s_{\star,1} \mathring{\mathcal{A}}_{\star,6} = 0. \quad (\text{B.32})$$

In the previous subsection we have shown by explicit computation that (B.28) and (B.29) imply $\mathring{\mathcal{A}}_{\star,3} = s_{\star,0} \mathcal{B}_3$ and $\mathring{\mathcal{A}}_{\star,4} = s_{\star,0} \mathcal{B}_4 + s_{\star,1} \mathcal{B}_3$. We shall now show by cohomological means that the remaining equations imply $\mathring{\mathcal{A}}_{\star,5} = s_{\star,0} \mathcal{B}_5 + s_{\star,1} \mathcal{B}_4$, $\mathring{\mathcal{A}}_{\star,6} = s_{\star,0} \mathcal{B}_6 + s_{\star,1} \mathcal{B}_5$ and $s_{\star,1} \mathcal{B}_6 = 0$, which will complete the proof of equations (B.9) to (B.13).

To that end we first derive a result on the cohomology of $s_{\star,0}$ in the space $\mathcal{F}_\star$ of integrated $\star$ -polynomials in the fields V_μ^A , Λ^A and their derivatives. An element of this space is a linear combination, with coefficients that may depend on $\omega^{\alpha\beta}$, of terms of the form

$$\int d^4x \, a_1 \star a_2 \star \dots \star a_n,$$

with n finite and each a_i one of our basic variables (V_μ^A , Λ^A and their derivatives),

$$a_i \in \{V_\mu^A, \Lambda^A, \partial_\mu V_\nu^A, \partial_\mu \Lambda^A, \partial_\mu \partial_\nu V_\rho^A, \partial_\mu \partial_\nu \Lambda^A, \dots\}.$$

It is obvious why this cohomology is relevant to the present case. Using the result $\mathring{\mathcal{A}}_{\star,4} = s_{\star,0} \mathcal{B}_4 + s_{\star,1} \mathcal{B}_3$ from Subappendix B.1 in eq. (B.30) and noting that $s_{\star,1}^2 = 0$, we obtain

$s_{\star,0}(\overset{\circ}{\mathcal{A}}_{\star,5} - s_{\star,1}\mathcal{B}_4) = 0$, with $\overset{\circ}{\mathcal{A}}_{\star,5} - s_{\star,1}\mathcal{B}_4$ obviously in $\mathcal{F}_\star$. Our aim is to show that this implies $\overset{\circ}{\mathcal{A}}_{\star,5} - s_{\star,1}\mathcal{B}_4 = s_{\star,0}\mathcal{B}_5$ for some $\mathcal{B}_5 \in \mathcal{F}_\star$, or in other words that $\overset{\circ}{\mathcal{A}}_{\star,5} - s_{\star,1}\mathcal{B}_4$ is trivial in the $s_{\star,0}$ -cohomology in $\mathcal{F}_\star$. Assume that we have shown this. Inserting the result in (B.31) and proceeding similarly yields $s_{\star,0}(\overset{\circ}{\mathcal{A}}_{\star,6} - s_{\star,1}\mathcal{B}_5) = 0$. Again, we want to show that $\overset{\circ}{\mathcal{A}}_{\star,6} - s_{\star,1}\mathcal{B}_5 = s_{\star,0}\mathcal{B}_6$ for some $\mathcal{B}_6 \in \mathcal{F}_\star$ and thus that $\overset{\circ}{\mathcal{A}}_{\star,6} - s_{\star,1}\mathcal{B}_5$ is also trivial in the $s_{\star,0}$ -cohomology in $\mathcal{F}_\star$. Note that (B.9) and (B.10) actually express analogous results, namely the triviality of $\overset{\circ}{\mathcal{A}}_{\star,3}$ and $\overset{\circ}{\mathcal{A}}_{\star,4} - s_{\star,1}\mathcal{B}_3$ in the same cohomology. However, as it will become clear below, they cannot be proved by means of the result on the cohomology for $s_{\star,0}$ in $\mathcal{F}_\star$ that we derive in the sequel and therefore have to be shown by other methods.

To examine the $s_{\star,0}$ -cohomology in $\mathcal{F}_\star$ we adapt methods developed in ref. [28] for the computation of the cohomology of s_0 . We first derive a result on the $s_{\star,0}$ -cohomology in the space $\mathcal{P}_\star$ of non-integrated $\star$ -polynomials. For that purpose we introduce the following variables u^ℓ , v^ℓ and w^i :

$$\{u^\ell\} = \{V_\mu^A, \partial_{(\mu} V_{\nu)}^A, \dots, \partial_{(\mu_1} \dots \partial_{\mu_k} V_{\mu_{k+1})}^A, \dots\} \quad (\text{B.33})$$

$$\{v^\ell\} = \{s_{\star,0}u^\ell\} = \{\partial_\mu \Lambda^A, \partial_{(\mu} \partial_{\nu)} \Lambda^A, \dots, \partial_{(\mu_1} \dots \partial_{\mu_{k+1})} \Lambda^A, \dots\} \quad (\text{B.34})$$

$$\{w^i\} = \{\Lambda^A, \partial_{[\mu} V_{\nu]}^A, \dots, \partial_{\mu_1} \dots \partial_{\mu_k} \partial_{[\mu} V_{\nu]}^A, \dots\}. \quad (\text{B.35})$$

Evidently every $\star$ -polynomial in the fields V_μ^A , Λ^A and their derivatives can be expressed as a $\star$ -polynomial in the variables u^ℓ , v^ℓ , w^i and vice versa⁴. On non-integrated $\star$ -monomials in u^ℓ , v^ℓ , w^i we define the operation ϱ through

$$\begin{aligned} \varrho(\hat{a}_1 \star \hat{a}_2 \star \dots \star \hat{a}_n) &= \\ &= \frac{1}{n} \left(u^\ell \frac{\partial \hat{a}_1}{\partial v^\ell} \right) \star \hat{a}_2 \star \dots \star \hat{a}_n \\ &+ \frac{1}{n} \sum_{i=2}^{n-1} (-)^{|\hat{a}_1|+|\hat{a}_2|+\dots+|\hat{a}_{i-1}|} \hat{a}_1 \star \dots \star \hat{a}_{i-1} \star \left(u^\ell \frac{\partial \hat{a}_i}{\partial v^\ell} \right) \star \hat{a}_{i+1} \star \dots \star \hat{a}_n \\ &+ \frac{1}{n} (-)^{|\hat{a}_1|+|\hat{a}_2|+\dots+|\hat{a}_{n-1}|} \hat{a}_1 \star \hat{a}_2 \star \dots \star \hat{a}_{n-1} \star \left(u^\ell \frac{\partial \hat{a}_n}{\partial v^\ell} \right), \end{aligned}$$

where $\hat{a}_i$ is any of the variables u^ℓ , v^ℓ , w^i ,

$$\hat{a}_i \in \{u^\ell, v^\ell, w^i\},$$

⁴The set of w 's is actually overcomplete because the w 's are not all linearly independent owing to the identities $\partial_{[\mu} \partial_{\nu]} V_{\rho]} = 0$ and their derivatives. However this does not matter to our arguments.

and $|\hat{a}_i|$ is the Grassmann parity of $\hat{a}_i$, which is 0 for V_μ^A and its derivatives, and 1 for the Λ^A and its derivatives. Extending the definition of ϱ by linearity from $\star$ -monomials to $\star$ -polynomials, we have that the anticommutator of $s_{\star,0}$ and ϱ evaluated on an arbitrary $\star$ -polynomial $p_\star(u, v, w) \in \mathcal{P}_\star$ gives the difference

$$\{s_{\star,0}, \varrho\} p_\star(u, v, w) = p_\star(u, v, w) - p_\star(0, 0, w) , \quad (\text{B.36})$$

where $p_\star(0, 0, w)$ denotes the $\star$ -polynomial that arises from $p_\star(u, v, w)$ by setting to zero all u^ℓ and v^ℓ *before* evaluating the star-products –for example, for $p_\star = V_\mu^A \star V_\nu^B$ one has $p_\star(0, 0, w) = 0$. Applying now eq. (B.36) to an $s_{\star,0}$ -closed $\star$ -polynomial, *i.e.* to a $p_\star$ satisfying $s_{\star,0}p_\star = 0$, and using that all w^i are $s_{\star,0}$ -closed, we obtain

$$s_{\star,0}p_\star(u, v, w) = 0 \quad \Leftrightarrow \quad p_\star(u, v, w) = p_\star(0, 0, w) + s_{\star,0}\varrho p_\star(u, v, w) . \quad (\text{B.37})$$

In particular, an $s_{\star,0}$ -closed $\star$ -polynomial $p_\star(u, v, w)$ with $p_\star(0, 0, w) = 0$ is the $s_{\star,0}$ -variation of the star-polynomial $\varrho p_\star(u, v, w)$.

Result (B.37) cannot be used directly for our purposes since it applies only to $\star$ -polynomials but not to integrated $\star$ -polynomials, which is what we had initially. This makes a difference because, by definition, an integrated $\star$ -polynomial is $s_{\star,0}$ -closed when the $s_{\star,0}$ -transformation of its integrand is a total divergence:

$$s_{\star,0}f_\star = 0 \quad \text{with} \quad f_\star = \int d^4x p_\star \quad \Leftrightarrow \quad s_{\star,0}p_\star = \partial_\mu \omega^\mu \quad \text{for some} \quad \omega^\mu .$$

Since ϱ does not commute with ∂_μ we cannot directly apply the result above to this case. To escape this problem we consider the variational derivatives of the equation $s_{\star,0}f_\star = 0$ with respect to V_μ^A and Λ^A . This yields

$$s_{\star,0}f_\star = 0, \quad f_\star \in \mathcal{F}_\star \quad \Rightarrow \quad s_{\star,0}\frac{\delta f_\star}{\delta V_\mu^A} = 0, \quad s_{\star,0}\frac{\delta f_\star}{\delta \Lambda^A} + \partial_\mu \frac{\delta f_\star}{\delta V_\mu^A} = 0 . \quad (\text{B.38})$$

It can be readily checked that the variational derivative of any element $f_\star \in \mathcal{F}_\star$ with respect to V_μ^A or Λ^A is a $\star$ -polynomial in $\mathcal{P}_\star$. Suppose now that $\delta f_\star/\delta V_\mu^A$ vanishes at $u^\ell = v^\ell = 0$ in the sense explained above. Using the first equation in (B.38) and eq. (B.37) we then conclude that $\delta f_\star/\delta V_\mu^A$ is the $s_{\star,0}$ -variation of $\varrho(\delta f_\star/\delta V_\mu^A)$:

$$\left[\frac{\delta f_\star}{\delta V_\mu^A} \right](0, 0, w) = 0 \quad \Rightarrow \quad \frac{\delta f_\star}{\delta V_\mu^A} = s_{\star,0}\varrho \frac{\delta f_\star}{\delta V_\mu^A} .$$

Using this in the second equation in (B.38) we obtain

$$s_{\star,0} \left(\frac{\delta f_{\star}}{\delta \Lambda^A} + \partial_{\mu} \varrho \frac{\delta f_{\star}}{\delta V_{\mu}^A} \right) = 0. \quad (\text{B.39})$$

Applying (B.37) once again we conclude that the term in parentheses is $s_{\star,0} \varrho(\dots)$ provided it vanishes at $u^{\ell} = v^{\ell} = 0$ in the sense above. Note that here $\varrho(\dots)$ has ghost number $\text{gh}(f_{\star}) - 2$, with $\text{gh}(f_{\star})$ the ghost number of $f_{\star}$ and $\text{gh}(V) = 0$ and $\text{gh}(\Lambda) = 1$. Since $\star$ -polynomials $p_{\star}(u, v, w)$ have non-negative ghost numbers, $\varrho(\dots)$ vanishes when $f_{\star}$ has ghost number 1, which is the case we are interested in. We thus conclude that

$$\left[\frac{\delta f_{\star}}{\delta \Lambda^A} + \partial_{\mu} \varrho \frac{\delta f_{\star}}{\delta V_{\mu}^A} \right] (0, 0, w) = 0, \quad \text{gh}(f_{\star}) = 1 \quad \Rightarrow \quad \frac{\delta f_{\star}}{\delta \Lambda^A} = -\partial_{\mu} \varrho \frac{\delta f_{\star}}{\delta V_{\mu}^A}. \quad (\text{B.40})$$

Finally we reconstruct $f_{\star}$ from its variational derivatives, neglecting integrated divergences, using the general formula

$$f_{\star}[V, \Lambda] = \int d^4x \int_0^1 \frac{d\tau}{\tau} \left(V_{\mu}^A \star \frac{\delta f_{\star}}{\delta V_{\mu}^A} + \Lambda^A \star \frac{\delta f_{\star}}{\delta \Lambda^A} \right) [\tau V, \tau \Lambda], \quad (\text{B.41})$$

valid for every functional $f_{\star}$. Using eqs. (B.39) and (B.40) in (B.41) we obtain

$$\begin{aligned} f_{\star}[V, \Lambda] &= \int d^4x \int_0^1 \frac{d\tau}{\tau} \left(V_{\mu}^A \star s_{\star,0} \varrho \frac{\delta f_{\star}}{\delta V_{\mu}^A} - \Lambda^A \star \partial_{\mu} \varrho \frac{\delta f_{\star}}{\delta V_{\mu}^A} \right) [\tau V, \tau \Lambda] \\ &= \int d^4x \int_0^1 \frac{d\tau}{\tau} \left(V_{\mu}^A \star s_{\star,0} \varrho \frac{\delta f_{\star}}{\delta V_{\mu}^A} + (s_{\star,0} V_{\mu}^A) \star \varrho \frac{\delta f_{\star}}{\delta V_{\mu}^A} \right) [\tau V, \tau \Lambda] \\ &= s_{\star,0} \int d^4x \int_0^1 \frac{d\tau}{\tau} \left(V_{\mu}^A \star \varrho \frac{\delta f_{\star}}{\delta V_{\mu}^A} \right) [\tau V, \tau \Lambda], \end{aligned}$$

where we have used integration by parts and $s_{\star,0} V_{\mu}^A = \partial_{\mu} \Lambda^A$. We have thus shown that

$$\begin{aligned} s_{\star,0} f_{\star} &= 0, \quad \text{gh}(f_{\star}) = 1, \quad \left[\frac{\delta f_{\star}}{\delta V_{\mu}^A} \right] (0, 0, w) = \left[\frac{\delta f_{\star}}{\delta \Lambda^A} + \partial_{\mu} \varrho \frac{\delta f_{\star}}{\delta V_{\mu}^A} \right] (0, 0, w) = 0 \\ \Rightarrow \quad f_{\star} &= s_{\star,0} \int d^4x \int_0^1 \frac{d\tau}{\tau} \left(V_{\mu}^A \star \varrho \frac{\delta f_{\star}}{\delta V_{\mu}^A} \right) [\tau V, \tau \Lambda], \end{aligned} \quad (\text{B.42})$$

which is the result for the $s_{\star,0}$ -cohomology in $\mathcal{F}_{\star}$ we will use to prove eqs. (B.30)-(B.32).

Consider now $\mathring{\mathcal{A}}_{\star,5} - s_{\star,1} \mathcal{B}_4$. It is an $s_{\star,0}$ -closed integrated $\star$ -polynomial with ghost number 1 whose integrand is order 5 in the fields V_{μ}^A and Λ^A , has mass dimension 4 –recall that

$\dim(V_\mu) = \dim(\partial_\mu) = 1$, $\dim(\Lambda) = 0$, $\dim(\omega^{\alpha\beta}) = -2$ – and contains one explicit $\omega^{\alpha\beta}$. It follows that the integrand is a linear combination of $\star$ -monomials $\omega^{\alpha\beta} a_1 \star \dots \star a_5$, where it can be assumed that one of the a_i is an undifferentiated Λ (for one can remove all derivatives from Λ using integrations by parts, if necessary) while the remaining a_i 's are either of type $\{V, V, \partial V, \partial V\}$ or $\{V, V, V, \partial \partial V\}$. It is easy to verify that this in turn implies

$$\left[\frac{\delta(\overset{\circ}{\mathcal{A}}_{\star,5} - s_{\star,1} \mathcal{B}_4)}{\delta V_\mu^A} \right] (0, 0, w) = 0 \quad (\text{B.43})$$

$$\left[\frac{\delta(\overset{\circ}{\mathcal{A}}_{\star,5} - s_{\star,1} \mathcal{B}_4)}{\delta \Lambda^A} + \partial_\mu \varrho \frac{\delta(\overset{\circ}{\mathcal{A}}_{\star,5} - s_{\star,1} \mathcal{B}_4)}{\delta V_\mu^A} \right] (0, 0, w) = 0. \quad (\text{B.44})$$

Eq. (B.42) can then be used and yields $\overset{\circ}{\mathcal{A}}_{\star,5} - s_{\star,1} \mathcal{B}_4 = s_{\star,0} \mathcal{B}_5$, with

$$\mathcal{B}_5 = \int d^4x \int_0^1 \frac{d\tau}{\tau} \left(V_\mu^A \star \varrho \frac{\delta(\overset{\circ}{\mathcal{A}}_{\star,5} - s_{\star,1} \mathcal{B}_4)}{\delta V_\mu^A} \right) [\tau V, \tau \Lambda].$$

This proves eq. (B.11) for $\mathcal{U}$ -valued fields, hence for $\mathfrak{g}$ -valued fields, as we wanted to show. The functional $\overset{\circ}{\mathcal{A}}_{\star,6} - s_{\star,1} \mathcal{B}_5$ can be treated analogously. Its integrand is order 6 in the fields, has mass dimension 4, ghost number 1 and one explicit $\omega^{\alpha\beta}$. It is thus a linear combination of $\star$ -monomials $\omega^{\alpha\beta} a_1 \star \dots \star a_6$, where it can be assumed that the set of a_i has the structure $\{\Lambda, V, V, V, V, \partial V\}$. This makes it obvious that $\overset{\circ}{\mathcal{A}}_{\star,6} - s_{\star,1} \mathcal{B}_5$ satisfies

$$\left[\frac{\delta(\overset{\circ}{\mathcal{A}}_{\star,6} - s_{\star,1} \mathcal{B}_5)}{\delta V_\mu^A} \right] (0, 0, w) = 0 \quad (\text{B.45})$$

$$\left[\frac{\delta(\overset{\circ}{\mathcal{A}}_{\star,6} - s_{\star,1} \mathcal{B}_5)}{\delta \Lambda^A} + \partial_\mu \varrho \frac{\delta(\overset{\circ}{\mathcal{A}}_{\star,6} - s_{\star,1} \mathcal{B}_5)}{\delta V_\mu^A} \right] (0, 0, w) = 0. \quad (\text{B.46})$$

Eq. (B.42) then implies $\overset{\circ}{\mathcal{A}}_{\star,6} - s_{\star,1} \mathcal{B}_5 = s_{\star,0} \mathcal{B}_6$, with

$$\mathcal{B}_6 = \int d^4x \int_0^1 \frac{d\tau}{\tau} \left(V_\mu^A \star \varrho \frac{\delta(\overset{\circ}{\mathcal{A}}_{\star,6} - s_{\star,1} \mathcal{B}_5)}{\delta V_\mu^A} \right) [\tau V, \tau \Lambda],$$

which proves eq. (B.12). Finally we have to show that eq. (B.13) holds. This is very easy. The integrand of $\mathcal{B}_6$ is a $\star$ -polynomial of order 6 in the fields, has mass dimension 4, ghost number 0 and one explicit $\omega^{\alpha\beta}$. It is thus a linear combination of $\star$ -monomials $\omega^{\alpha\beta} a_1 \star \dots \star a_6$, where all a_i are undifferentiated V 's. Furthermore, by construction, it can be written as a trace Tr . The latter implies already $s_{\star,1} \mathcal{B}_6 = 0$, since

$$s_{\star,1} \text{Tr} \left(V_{\mu_1} \star \dots \star V_{\mu_6} \right) = \text{Tr} \left[V_{\mu_1} \star \dots \star V_{\mu_6}, \Lambda \right]_\star$$

is a divergence.

We close by remarking that eq. (B.42) cannot be used to prove that $\mathring{\mathcal{A}}_{\star,3}$ and $\mathring{\mathcal{A}}_{\star,4} - s_{\star,1}\mathcal{B}_3$ are trivial in the $s_{\star,0}$ -cohomology in $\mathcal{F}_\star$ because the $\frac{\delta}{\delta V_\mu^A}$ and $\frac{\delta}{\delta \Lambda^A} + \partial_\mu e \frac{\delta}{\delta V_\mu^A}$ acting on them do not vanish at $u^\ell = v^\ell = 0$ in the sense explained above, contrary to what happens for $\mathring{\mathcal{A}}_{\star,5} - s_{\star,1}\mathcal{B}_4$ and $\mathring{\mathcal{A}}_{\star,6} - s_{\star,1}\mathcal{B}_5$ –see eqs. (B.43), (B.44), (B.45) and (B.46).

References

- [1] J. Madore, S. Schraml, P. Schupp and J. Wess, “Gauge theory on noncommutative spaces,” Eur. Phys. J. C **16** (2000) 161 [arXiv:hep-th/0001203].
- [2] B. Jurco, S. Schraml, P. Schupp and J. Wess, “Enveloping algebra valued gauge transformations for non-Abelian gauge groups on non-commutative spaces,” Eur. Phys. J. C **17** (2000) 521 [arXiv:hep-th/0006246].
- [3] N. Seiberg and E. Witten, “String theory and noncommutative geometry,” JHEP **9909** (1999) 032 [arXiv:hep-th/9908142].
- [4] B. Jurco, P. Schupp and J. Wess, “Nonabelian noncommutative gauge theory via non-commutative extra dimensions,” Nucl. Phys. B **604** (2001) 148 [arXiv:hep-th/0102129].
- [5] B. Jurco, L. Moller, S. Schraml, P. Schupp and J. Wess, “Construction of non-Abelian gauge theories on noncommutative spaces,” Eur. Phys. J. C **21** (2001) 383 [arXiv:hep-th/0104153].
- [6] X. Calmet, B. Jurco, P. Schupp, J. Wess and M. Wohlgenannt, “The standard model on non-commutative space-time,” Eur. Phys. J. C **23** (2002) 363 [arXiv:hep-ph/0111115].
- [7] S. M. Carroll, J. A. Harvey, V. A. Kostelecky, C. D. Lane and T. Okamoto, “Non-commutative field theory and Lorentz violation,” Phys. Rev. Lett. **87** (2001) 141601 [arXiv:hep-th/0105082].
- [8] C. E. Carlson, C. D. Carone and R. F. Lebed, “Bounding noncommutative QCD,” Phys. Lett. B **518** (2001) 201 [arXiv:hep-ph/0107291].
- [9] W. Behr, N. G. Deshpande, G. Duplancic, P. Schupp, J. Trampetic and J. Wess, “The $Z \rightarrow \gamma\gamma, gg$ decays in the noncommutative standard model,” [arXiv:hep-ph/0202121].

- [10] C. E. Carlson, C. D. Carone and R. F. Lebed, “Supersymmetric noncommutative QED and Lorentz violation,” *Phys. Lett. B* **549** (2002) 337 [arXiv:hep-ph/0209077].
- [11] P. Schupp, J. Trampetic, J. Wess and G. Raffelt, “The photon neutrino interaction in non-commutative gauge field theory and astrophysical bounds,” arXiv:hep-ph/0212292.
- [12] P. Minkowski, P. Schupp and J. Trampetic, “Non-commutative ‘*-charge radius’ and ‘*-dipole moment’ of the neutrino,” arXiv:hep-th/0302175.
- [13] P. Aschieri, B. Jurco, P. Schupp and J. Wess, “Non-commutative GUTs, standard model and C, P, T,” *Nucl. Phys. B* **651** (2003) 45 [arXiv:hep-th/0205214].
- [14] G. Barnich and M. Henneaux, “Renormalization of gauge invariant operators and anomalies in Yang-Mills theory,” *Phys. Rev. Lett.* **72** (1994) 1588 [arXiv:hep-th/9312206].
- [15] G. Barnich, F. Brandt and M. Henneaux, “Local BRST cohomology in the antifield formalism: II. Application to Yang-Mills theory,” *Commun. Math. Phys.* **174** (1995) 93 [arXiv:hep-th/9405194].
- [16] G. Barnich, F. Brandt and M. Henneaux, “Local BRST cohomology in gauge theories,” *Phys. Rept.* **338** (2000) 439 [arXiv:hep-th/0002245].
- [17] R. Wulkenhaar, “Non-renormalizability of theta-expanded noncommutative QED,” *JHEP* **0203** (2002) 024 [arXiv:hep-th/0112248].
- [18] C. P. Martin, “The gauge anomaly and the Seiberg-Witten map,” *Nucl. Phys. B* **652** (2003) 72 [arXiv:hep-th/0211164].
- [19] R. Banerjee and S. Ghosh, “Seiberg-Witten map and the axial anomaly in noncommutative field theory,” *Phys. Lett. B* **533** (2002) 162 [arXiv:hep-th/0110177].
- [20] R. Banerjee, “Anomalies in noncommutative gauge theories, Seiberg-Witten transformation and Ramond-Ramond couplings,” arXiv:hep-th/0301174.
- [21] B. L. Cerchiai, A. F. Pasqua and B. Zumino, arXiv:hep-th/0206231.
- [22] P. Breitenlohner and D. Maison, “Dimensional renormalization and the action principle,” *Commun. Math. Phys.* **52** (1977) 11.

- [23] G. Barnich, F. Brandt and M. Grigoriev, “Seiberg-Witten maps and noncommutative Yang-Mills theories for arbitrary gauge groups,” JHEP **0208**, 023 (2002) [arXiv:hep-th/0206003].
- [24] H. Georgi and S. L. Glashow, “Gauge Theories Without Anomalies,” Phys. Rev. D **6** (1972) 429.
- [25] J. M. Gracia-Bondia and C. P. Martin, “Chiral gauge anomalies on noncommutative R^{*4} ,” Phys. Lett. B **479** (2000) 321 [arXiv:hep-th/0002171].
- [26] L. Bonora, M. Schnabl and A. Tomasiello, “A note on consistent anomalies in noncommutative YM theories,” Phys. Lett. B **485** (2000) 311 [arXiv:hep-th/0002210].
- [27] D. J. Gross and R. Jackiw, “Effect Of Anomalies On Quasirenormalizable Theories,” Phys. Rev. D **6** (1972) 477.
- [28] F. Brandt, N. Dragon and M. Kreuzer, “Completeness and nontriviality of the solutions of the consistency conditions,” Nucl. Phys. B **332** (1990) 224.